

## Overview

### Useful For

Screening for liver damage, especially if someone has a condition or is taking a drug that may affect the liver

### Method Name

BILIT: Photometric, Diazonium Salt

BILID: Photometric, Diazotized Sulfanilic Acid

AST: Photometric Rate, L-Aspartate with Pyridoxyl-5-Phosphate

ALT: Photometric Rate, L-Alanine with Pyridoxal-5-Phosphate

ALP: Photometric, p-Nitrophenol Phosphate

ALB: Photometric, Bromocresol Green

TP: Colorimetric, Biuret

### NY State Available

Yes

## Specimen

### Specimen Type

Serum

### Shipping Instructions

[Ship specimen in amber vial to protect from light.](#)

### Necessary Information

Patient's age and sex are required.

### Specimen Required

**Supplies:** Amber Frosted Tube, 5 mL (T915)

**Collection Container/Tube:**

**Preferred:** Serum gel

**Acceptable:** Red top

**Submission Container/Tube:** Amber vial

**Specimen Volume:** 0.6 mL

**Collection Instructions:**

1. Serum gel tubes should be centrifuged within 2 hours of collection.
2. Red-top tubes should be centrifuged, and the serum aliquoted into an amber vial within 2 hours of collection.

### Specimen Minimum Volume

0.5 mL

**Reject Due To**

Gross hemolysis	Reject
Gross lipemia	OK

**Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated	7 days	

**Clinical & Interpretive**
**Clinical Information**

The hepatic function panel may be used to help diagnose liver disease if a person has signs and symptoms that indicate possible liver dysfunction. If a person has a known condition or liver disease, testing may be performed at intervals to monitor the health of the liver and to evaluate the effectiveness of any treatments. Abnormal tests on a liver panel may prompt a repeat analysis of one or more tests, or of the whole panel, to see if the elevations or decreases persist and may indicate the need for additional testing to determine the cause of the liver dysfunction.

**Reference Values**
**TOTAL BILIRUBIN**

0-6 days: Refer to <http://bilitool.org/> for information on age-specific (postnatal hour of life) serum bilirubin values.

7-14 days: 0.0-14.9

15 days to 17 years: 0.0 -1.0

>18 years: 0.0-1.2

**DIRECT BILIRUBIN**

> or =12 months: 0.0-0.3 mg/dL

Reference values have not been established for patients who are younger than 12 months of age.

**ASPARTATE AMINOTRANSFERASE**
**Males**

0-11 months: Not established

1-13 years: 8-60 U/L

> or =14 years: 8-48 U/L

**Females**

0-11 months: Not established

1-13 years: 8-50 U/L

> or =14 years: 8-43 U/L

**ALANINE AMINOTRANSFERASE**
**Males**

---

> or =1 year: 7-55 U/L

Reference values have not been established for patients who are younger than 12 months of age.

Females

> or =1 year: 7-45 U/L

Reference values have not been established for patients who are younger than 12 months of age.

#### ALKALINE PHOSPHATASE

Males

4 years: 149-369 U/L

5 years: 179-416 U/L

6 years: 179-417 U/L

7 years: 172-405 U/L

8 years: 169-401 U/L

9 years: 175-411 U/L

10 years: 191-435 U/L

11 years: 185-507 U/L

12 years: 185-562 U/L

13 years: 182-587 U/L

14 years: 166-571 U/L

15 years: 138-511 U/L

16 years: 102-417 U/L

17 years: 69-311 U/L

18 years: 52-222 U/L

> or =19 years: 45-115 U/L

Females

4 years: 169-372 U/L

5 years: 162-355 U/L

6 years: 169-370 U/L

7 years: 183-402 U/L

8 years: 199-440 U/L

9 years: 212-468 U/L

10 years: 215-476 U/L

11 years: 178-526 U/L

12 years: 133-485 U/L

13 years: 120-449 U/L

14 years: 153-362 U/L

15 years: 75-274 U/L

16 years: 61-264 U/L

17-23 years: 52-144 U/L

24-45 years: 37-98 U/L

46-50 years: 39-100 U/L

51-55 years: 41-108 U/L

56-60 years: 46-118 U/L

---

61-65 years: 50-130 U/L

> or =66 years: 55-142 U/L

Reference values have not been established for patients who are younger than 4 years of age.

**ALBUMIN**

> or =12 months: 3.5-5.0 g/dL

Reference values have not been established for patients who are younger than 12 months of age.

**TOTAL PROTEIN**

> or =1 year: 6.3-7.9 g/dL

Reference values have not been established for patients who are younger than 12 months of age.

**Interpretation**

Hepatic function panel results are not diagnostic of a specific condition; they indicate that there may be a problem with the liver. In a person who does not have symptoms or identifiable risk factors, abnormal liver test results may indicate a temporary liver injury or reflect something that is happening elsewhere in the body, such as in the skeletal muscles, pancreas, or heart. It may also indicate early liver disease and the need for further testing and periodic monitoring.

Results of liver panels are usually evaluated together. Several sets of results from tests performed over a few days or weeks are often assessed together to determine if a pattern is present. Each person will have a unique set of test results that will typically change over time. A healthcare practitioner evaluates the combination of liver test results to gain clues about the underlying condition. Often, additional testing is necessary to determine what is causing the liver damage or disease.

**Cautions**

No significant cautionary statements

**Clinical Reference**

Kampfrath T. Liver Panel Test. Testing.com; Updated September 28, 2022. Accessed September 19, 2023. Available at <https://www.testing.com/tests/liver-panel/>

**Performance****Method Description**

Bilirubin, Total:

Total bilirubin, in the presence of a suitable solubilizing agent, is coupled with 3,5-dichlorophenyl diazonium in a strongly acidic medium. The color intensity of the red azo dye formed is directly proportional to the total bilirubin and can be determined photometrically.(Package insert: Bilirubin Total Gen. 3, 07/2014. Roche Diagnostics, Indianapolis, IN)

Bilirubin, Direct:

Acidified sodium nitrite produces nitrous acid which reacts with sulfanilic acid (in acidic solution) to form a diazonium salt. The diazotized sulfanilic acid then reacts with bilirubin to form isomers of azobilirubin. In the direct bilirubin assay, only conjugated bilirubin is converted by the diazotized sulfanilic acid. The intensity of the color of azobilirubin is measured photometrically at 570 nm and is proportional to the direct (conjugated) bilirubin concentration.(Package

---

insert: Roche Direct Bilirubin reagent; Indianapolis, IN, October 1999)

**Aspartate Aminotransferase (AST):**

AST is measured by a coupled enzyme kinetic method where the rate of decrease of NADH, determined at 340 nm, is directly proportional to the AST activity. (Package insert: Roche AST reagent, Indianapolis, IN, January 2000)

**Alanine Aminotransferase (ALT):**

ALT activity is determined by a kinetic method using a coupled enzyme reaction where the rate of NADH consumption is measured at 340 nm. The NADH decrease is directly proportional to the ALT activity. (Package insert: Roche ALT reagent, Indianapolis, IN, January 2000)

**Alkaline Phosphatase:**

In the presence of magnesium and zinc ions, p-nitrophenyl phosphate is cleaved by phosphatases into phosphate and p-nitrophenol. The p-nitrophenol released is directly proportional to the catalytic alkaline phosphatase activity. It is determined by measuring the increase in absorbance. (Package insert: Roche Alkaline Phosphatase reagent, Indianapolis, IN, February 2012)

**Albumin:**

The dye, bromocresol green (BCG), is added to serum in an acid buffer. The color intensity of the blue-green albumin-BCG complex is directly proportional to the albumin concentration and is determined photometrically. (Package insert: Roche Albumin reagent; Roche Diagnostic Corp., Indianapolis, IN, July 1999)

**Protein, Total:**

Divalent copper reacts in alkaline solution with protein peptide bonds to form the characteristic purple-colored biuret complex. Sodium potassium tartrate prevents the precipitation of copper hydroxide and potassium iodide prevents autoreduction of copper. The color intensity is directly proportional to the protein concentration which can be determined photometrically. (Package insert: Roche Protein reagent, Roche Diagnostic Corp., Indianapolis, IN 1999)

**PDF Report**

No

**Day(s) Performed**

Monday through Sunday

**Report Available**

Same day/1 to 2 days

**Specimen Retention Time**

1 week

**Performing Laboratory Location**

Rochester

**Fees & Codes**

## Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

## Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

## CPT Code Information

- 82247
- 82248
- 84450
- 84460
- 84075
- 82040
- 84155
- 80076 (if appropriate if all analytes performed)

## LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
LIVPR	Hepatic Function Panel, S	24325-3

Result ID	Test Result Name	Result LOINC® Value
ALB	Albumin, S	1751-7
ALP	Alkaline Phosphatase, S	6768-6
ALT	Alanine Aminotransferase (ALT), S	1743-4
AST	Aspartate Aminotransferase (AST), S	30239-8
BILID	Bilirubin, Direct	1968-7
BILIT	Bilirubin Total, S	1975-2
TP	Protein, Total, S	2885-2