

Overview

Useful For

Identifying *MYC* amplification to aid in the differentiation of cutaneous angiosarcomas from atypical vascular lesions after radiotherapy

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
_PBCT	Probe, +2	No, (Bill Only)	No
_PADD	Probe, +1	No, (Bill Only)	No
_PB02	Probe, +2	No, (Bill Only)	No
_PB03	Probe, +3	No, (Bill Only)	No
_IL25	Interphases, <25	No, (Bill Only)	No
_I099	Interphases, 25-99	No, (Bill Only)	No
_I300	Interphases, >=100	No, (Bill Only)	No

Testing Algorithm

This test does not include a pathology consultation. If a pathology consult is requested, PATHC / Pathology Consultation should be ordered, and the appropriate fluorescence in situ hybridization (FISH) test will be performed at an additional charge.

This test includes a charge for application of the first probe set (2 FISH probes) and professional interpretation of results.

Additional charges will be incurred for all reflex probes performed. Analysis charges will be incurred based on the number of cells analyzed per probe set. If no cells are available for analysis, no analysis charges will be incurred.

Appropriate ancillary probes may be performed at consultant discretion to render comprehensive assessment. Any additional probes will have the results included within the final report and will be performed at an additional charge.

Method Name

Fluorescence In Situ Hybridization (FISH)

NY State Available

Yes

Specimen

Specimen Type

Tissue

Ordering Guidance

This test does not include a pathology consultation. If a pathology consultation is requested, order PATHC / Pathology Consultation, and appropriate testing will be added at the discretion of the pathologist and performed at an additional charge.

Multiple oncology (cancer) gene panels are also available. For more information see [Hematology, Oncology, and Hereditary Test Selection Guide](#).

Additional Testing Requirements

To resolve atypical fluorescence in situ hybridization results, confirmation testing by microarray testing is available; order CMAPT / Chromosomal Microarray, Tumor, Formalin-Fixed Paraffin-Embedded.

Shipping Instructions

Advise Express Mail or equivalent if not on courier service.

Necessary Information

1. A pathology report is required for testing to be performed. If not provided, appropriate testing and interpretation may be compromised or delayed. Acceptable pathology reports include working drafts, preliminary pathology, or surgical pathology reports.

2. The following information must be included in the report provided:

- Patient name
- Block number - must be on all blocks, slides, and paperwork
- Date of collection
- Tissue source

3. A reason for testing must be provided. If this information is not provided, an appropriate indication for testing may be entered by Mayo Clinic Laboratories.

Specimen Required

Submit only 1 of the following specimens:

Preferred:

Specimen Type: Tissue block

Collection Instructions: Submit a formalin-fixed, paraffin-embedded tumor tissue block. Blocks prepared with alternative fixation methods will be attempted but are less favorable for successful results by fluorescence in situ hybridization testing; provide fixation method used.

Additional Information:

1. Paraffin-embedded specimens can be from any anatomic location (skin, soft tissue, lymph node, etc).
2. Bone specimens that have been decalcified will be attempted for testing, but the success rate is approximately 50%.

Acceptable:

Specimen Type: Tissue slides

Slides: 1 Hematoxylin and eosin stained and 4 unstained

Collection Instructions: Submit 1 slide stained with hematoxylin and eosin and 4 consecutive unstained, positively

charged, unbaked slides with 5 micron-thick sections of the tumor tissue.

Forms

If not ordering electronically, complete, print, and send an [Oncology Test Request](#) (T729) with the specimen.

Specimen Minimum Volume

Slides: 1 Hematoxylin and eosin stained and 2 unstained

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Tissue	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive**Clinical Information**

Postradiation cutaneous angiosarcoma is a malignancy associated with very poor outcome and is consequently treated aggressively. Atypical vascular lesions are also associated with radiation therapy but are considered to be benign and do not require aggressive management. Therefore, the differentiation of these neoplasms is of considerable clinical importance. Postradiation cutaneous angiosarcomas often demonstrate high-level amplification of *MYC*, whereas reactive and benign vascular lesions do not show amplification of *MYC*.

Reference Values

An interpretive report will be provided.

Interpretation

MYC will be clinically interpreted as positive, negative, or equivocal.

The *MYC* locus is reported as amplified when the *MYC*:D8Z2 ratio is 2.0 or greater and demonstrates 6 or more copies of *MYC*.

A *MYC*:D8Z2 ratio less than 2.0 or showing a ratio of 2.0 or greater with less than 6 copies of *MYC* is considered to lack amplification of *MYC*.

Cautions

This test is not approved by the US Food and Drug Administration, and it is best used as an adjunct to existing clinical and pathologic information.

This fluorescence in situ hybridization (FISH) assay does not rule out other chromosome abnormalities.

Fixatives other than formalin (eg, Prefer, Bouin's) may not be successful for FISH assays. Non-formalin fixed specimens will not be rejected.

Paraffin-embedded tissues that have been decalcified may not be successful for FISH analysis. The success rate of FISH studies on decalcified tissue is approximately 50%, but FISH will be attempted if sufficient tumor is present for analysis.

FISH studies will be attempted if sufficient tumor is present for analysis. The pathologist reviewing the hematoxylin and eosin-stained slide may find it necessary to cancel testing if insufficient tissue/tumor is available for testing.

If no FISH signals are observed post hybridization, the case will be released indicating a lack of FISH results.

Supportive Data

The probe set was independently validated in a blinded study on 23 paraffin-embedded primary and post radiation angiosarcoma tissue samples and 25 noncancerous control specimens. The normal controls were used to generate the normal cutoffs. *MYC* amplification was detected in 4 (17.4%) of the angiosarcomas and the incidence is consistent with published reports.

Clinical Reference

1. Mentzel T, Schildhaus H, Palmedo G, Buttner R, Kutzner H. Postradiation cutaneous angiosarcoma after treatment of breast carcinoma is characterized by *MYC* amplification in contrast to atypical vascular lesions after radiotherapy and control cases: clinicopathological, immunohistochemical and molecular analysis of 66 cases. *Mod Pathol*. 2012;25(1):75-85
2. Manner J, Radlwimmer B, Hohenberger P, et al. *MYC* high level gene amplification is a distinctive feature of angiosarcomas after irradiation or chronic lymphedema. *Am J Pathol*. 2010;176(1):34-39
3. WHO Classification of Tumours Editorial Board. Soft Tissue and Bone. 5th ed. IARC; 2020. World Health Organization Classification of Tumours. Vol 3

Performance

Method Description

This test is performed using a commercially available MYC probe set with a MYC probe and a chromosome 8 centromere probe (D8Z2). The selection of tissue and the identification of target areas on the hematoxylin and eosin (H and E)-stained slide are performed by a pathologist. Using the H and E-stained slide as a reference, target areas are etched with a diamond-tipped engraving tool on the back of the unstained slide to be assayed. The probe set is hybridized to the appropriate target areas, and 2 technologists each independently analyze 30 interphase nuclei (60 total) with the results expressed as a ratio of MYC:D8Z2 signals.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

7 to 10 days

Specimen Retention Time

Slides and H and E used for analysis are retained by the laboratory in accordance with regulatory requirements. Client provided paraffin blocks and extra unstained slides (if provided) will be returned after testing is complete.

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

- 88271x2, 88291-DNA probe, each (first probe set), Interpretation and report
- 88271x2-DNA probe, each; each additional probe set (if appropriate)
- 88271x1-DNA probe, each; coverage for sets containing 3 probes (if appropriate)
- 88271x2-DNA probe, each; coverage for sets containing 4 probes (if appropriate)
- 88271x3-DNA probe, each; coverage for sets containing 5 probes (if appropriate)
- 88274 w/modifier 52-Interphase in situ hybridization, <25 cells, each probe set (if appropriate)
- 88274-Interphase in situ hybridization, 25 to 99 cells, each probe set (if appropriate)
- 88275-Interphase in situ hybridization, 100 to 300 cells, each probe set (if appropriate)

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
MASF	MYC (8q24), Angiosarcoma, FISH, Ts	101384-6

Result ID	Test Result Name	Result LOINC® Value
54606	Result Summary	50397-9
54609	Interpretation	69965-2
54608	Result	62356-1
CG896	Reason for Referral	42349-1
54610	Specimen	31208-2
54611	Source	31208-2

Test Definition: MASF

Angiosarcoma, MYC (8q24) Amplification,
FISH, Tissue

54612	Tissue ID	80398-1
54613	Method	85069-3
54614	Released By	18771-6
55126	Additional Information	48767-8
53818	Disclaimer	62364-5