

## Overview

### Useful For

Detecting the presence or helping to exclude the presence of lupus anticoagulants (LA)

Identifying LA that do not prolong the activated partial thromboplastin time (APTT)

Evaluating unexplained prolongation of the APTT or prothrombin time clotting tests

Distinguishing LA from a specific coagulation factor inhibitor or coagulation factor deficiencies

Confirmation of abnormal dilute Russell's viper venom time ratios

### Method Name

Only orderable as part of a reflex. For more information see:

ALUPP / Lupus Anticoagulant Profile, Plasma

ALBLD / Bleeding Diathesis Profile, Limited, Plasma

AATHR / Thrombophilia Profile, Plasma and Whole Blood

APROL / Prolonged Clot Time Profile, Plasma

ADIC / Disseminated Intravascular Coagulation/Intravascular Coagulation and Fibrinolysis (DIC/ICF) Profile, Plasma

Optical Clot-Based

### NY State Available

Yes

## Specimen

### Specimen Type

Plasma Na Cit

### Specimen Required

Only orderable as part of a reflex. For more information see:

ALUPP / Lupus Anticoagulant Profile, Plasma

ALBLD / Bleeding Diathesis Profile, Limited, Plasma

AATHR / Thrombophilia Profile, Plasma and Whole Blood

APROL / Prolonged Clot Time Profile, Plasma

ADIC / Disseminated Intravascular Coagulation/Intravascular Coagulation and Fibrinolysis (DIC/ICF) Profile, Plasma

### Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject
Gross icterus	Reject

### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Plasma Na Cit	Frozen	14 days	

### Clinical & Interpretive

#### Clinical Information

Lupus anticoagulants (LA) are immunoglobulins (IgG, IgM, IgA, or a combination of these) of autoimmune type that are specifically directed against antigenic complexes of negatively charged phospholipids (such as phosphatidylserine or phosphatidylethanolamine) and coagulation-related proteins (such as beta-2-glycoprotein I or clotting factors including prothrombin [factor II] or factor X), and which cause prolongation of phospholipid-dependent clotting time tests due to inhibition.

LA are functionally and clinically distinct members of a broader group of antiphospholipid autoantibodies that includes immunologically detectable anticardiolipin antibodies or antibodies against other phospholipid-protein complexes. LA interfere with specific coagulation factor-phospholipid interactions, typically causing prolongation of 1 or more phospholipid-dependent clotting time tests (eg, activated partial thromboplastin time: APTT, dilute Russell's viper venom time: DRVVT) due to inhibition. This characteristic *in vitro* inhibition can be overcome by addition of excess phospholipid.

Because of the heterogeneous nature of LA antibodies, no single coagulation test can identify or exclude all LA. Currently, the International Society on Thrombosis and Haemostasis and the Clinical and Laboratory Standards Institute (CLSI) recommend testing for LA with at least 2 phospholipid-dependent clotting time assays based on different coagulation pathways and principles (eg, lupus-sensitive APTT and DRVVT).

In addition, given the potential for false-positive results in patients on anticoagulants, a profile or panel of coagulation tests is performed, including the prothrombin time (PT), APTT, thrombin time (TT) and DRVVT. If the PT, APTT, and/or DRVVT are prolonged, additional testing may include mixing tests with normal plasma (to evaluate for inhibition) and the use of excess phospholipid in appropriate assay systems to evaluate for phospholipid-dependent inhibition. Additional reflexive testing helps determine presence or absence of anticoagulants and/or inhibitors to other factors.

The diagnosis of LA requires performance and interpretation of complex coagulation testing, as well as correlation with available clinical information, including evidence of persistence of LA over time (> or =12 weeks).

The venom obtained from Russell's viper (*Vipera russelli*) contains enzymes that directly activate coagulation factors V

---

and X, bypassing the activation of factors VII, VIII, IX, XI, and XII, and, therefore, the effect of deficiencies or inhibitors of these factors. Diluting the phospholipid necessary for the clotting factor interactions increases the sensitivity to LA and the likelihood of identifying a phospholipid-dependent inhibitor that may not be detected by other coagulation tests with a higher phospholipid content (eg, LA-insensitive APTT reagents).

The DRVVT screen ratio test is one of several available in vitro tests that may be used to screen and confirm for presence of LA or to help exclude LA. DRVVT testing is used in conjunction with other appropriate coagulation tests (reflexive testing panels) to assist in detection and confirmation of LA, or help exclude their presence.

The DRVVT may be abnormally prolonged (DRVVT screen ratio  $>$  or  $\geq$ 1.20) by LA as well as coagulation factor deficiencies, anticoagulant effects, or other types of coagulation factor inhibitors.

Specimens with abnormal results (DRVVT screen ratio  $>$  or  $\geq$ 1.20) are subjected to reflexive testing. With the reflexive testing algorithm, the sensitivity of DRVVT testing for LA diagnosis is approximately 65% to 70% and the specificity is 95% or higher.

Although LA cause prolonged clotting times in vitro, there is a strong association with thrombosis risk. However, not all patients with persisting LA develop thrombosis.

## Reference Values

Only orderable as part of a reflex. For more information see:

ALUPP / Lupus Anticoagulant Profile, Plasma

ALBLD / Bleeding Diathesis Profile, Limited, Plasma

AATHR / Thrombophilia Profile, Plasma and Whole Blood

APROL / Prolonged Clot Time Profile, Plasma

ADIC / Disseminated Intravascular Coagulation/Intravascular Coagulation and Fibrinolysis (DIC/ICF) Profile, Plasma

<1.20

Normal ranges for children: not clearly established, but similar to normal ranges for adults, except for newborn infants whose results may not reach adult values until 3 to 6 months of age.

## Interpretation

Dilute Russell's viper venom time (DRVVT) screen ratio ( $<1.20$ ):

A normal DRVVT screen ratio ( $<1.20$ ) indicates that lupus anticoagulant (LA) is not present or not detectable by this method (but might be detected with other methods).

An abnormal DRVVT screen ratio (DRVVT screen ratio  $>1.20$ ) may suggest presence of LA, however, other possibilities include:

-Deficiencies or dysfunction of factors I (fibrinogen), II, V, or X, congenital or acquired

-Inhibitors of factor V, or occasionally by inhibitors of factor VIII, or other specific or nonspecific inhibitors

-Anticoagulation therapy effects (see Cautions)

Further evaluation consists of performing mixing studies with an equal volume of normal pooled plasma (DRVVT 1:1 mix) to investigate the possibility of coagulation factor deficiency (suggested by DRVVT mix ratio  $<1.20$ ) and to evaluate

---

inhibition (suggested by DRVVT mix ratio  $>$  or  $\geq$  1.20) and mixing patient plasma with DRVVT reagent enriched in phospholipid (DRVVT confirmatory reagent) (DRVVT mix and DRVVT confirmation ratios).

Possible combination of results include the following:

-DRVVT screen ratio  $>$  or  $\geq$  1.20, DRVVT mix ratio  $<$  1.20, and DRVVT confirmation ratio  $<$  1.20:

No evidence of LA. These data may reflect anticoagulation therapy effects or other (congenital or acquired) coagulopathy.

-DRVVT screen ratio  $>$  or  $\geq$  1.20, DRVVT mix ratio  $>$  or  $\geq$  1.20, and DRVVT confirmation ratio  $<$  1.20:

The prolonged and inhibited DRVVT (DRVVT screen and mix ratios) may reflect presence of a specific factor inhibitor (eg, factor V inhibitor), anticoagulation therapy effects or other nonspecific inhibitors as can be seen with monoclonal protein disorders, lymphoproliferative disease etc. Although LA cannot be conclusively excluded, the DRVVT confirmation ratio of  $<$  or  $\geq$  1.20 makes this less likely.

-DRVVT screen ratio  $>$  or  $\geq$  1.20, DRVVT mix ratio  $<$  1.20, and DRVVT confirmation ratio  $>$  or  $\geq$  1.20:

Although mixing study of the prolonged DRVVT screen and mix ratios provides no evidence of inhibition, additional phospholipid shortens the clotting time (DRVVT confirmation ratio), suggesting presence of LA.

-DRVVT screen ratio  $>$  or  $\geq$  1.20, DRVVT mix ratio  $>$  or  $\geq$  1.20, and DRVVT confirmation ratio  $>$  or  $\geq$  1.20:

The data are consistent with presence of LA, provided anticoagulant effect can be excluded (see Cautions)

Additional tests to evaluate abnormal DRVVT results include activated partial thromboplastin time (APTT), prothrombin time (PT), and thrombin time. Abnormalities observed with these tests may be further evaluated with normal plasma mixing studies, the platelet neutralization procedure (for APTT), and coagulation factor assays may sometimes be needed. All of these reflexive testing procedures, together with Coagulation Consultant interpretation, are included in Mayo Clinic's Coagulation Consultation test panels:

ALUPP / Lupus Anticoagulant Profile, Plasma

ALBLD / Bleeding Diathesis Profile, Limited, Plasma

AATHR / Thrombophilia Profile, Plasma and Whole Blood

APROL / Prolonged Clot Time Profile, Plasma

ADIC / Disseminated Intravascular Coagulation/Intravascular Coagulation and Fibrinolysis (DIC/ICF) Profile, Plasma

DRVVT assays ordered as a single, stand-alone test should be interpreted within patient clinical context and close attention to medication use by patient (see Cautions).

### **Cautions**

Residual platelets in frozen-thawed plasma can decrease sensitivity and specificity of lupus anticoagulants (LA) testing (false-negative results). Specimens that are to be frozen before testing must be centrifuged twice to remove as many of the platelets as possible before freezing.

Anticoagulation therapy effects such as warfarin (especially when the effect is supratherapeutic), excess heparin, direct thrombin inhibitors (eg, dabigatran [Pradaxa]), argatroban [Ancova], bivalirudin [Angiomax]), direct factor Xa inhibitors (eg, rivaroxaban [Xarelto], apixaban [Eliquis], edoxaban [Savaysa]) may result in a false-positive assay performance for LA. Clinical correlation and repeat testing remote ( $>1$  week) from anticoagulation therapy is suggested.

Although the dilute Russell's viper venom time (DRVVT) reagents contain a heparin inhibitor (Polybrene) that is sufficient for neutralization of heparin (up to 1-2 U/mL), the results may not necessarily represent what would occur if no heparin

were present in the specimen. Therefore, DRVVT results from heparinized plasma should be interpreted with caution.

Excess heparin or inhibitors of factors V or VIII may cause false-positive results of LA testing, depending on the types of coagulation testing performed.

LA diagnosis does not have definite predictive value for associated clinical complications such as thromboembolic problems or fetal loss.

The DRVVT test will not detect all LAs. Some LAs may only be detectable by other tests such as the Staclot LA, activated partial thromboplastin time, and platelet neutralization procedure, or other methods.

Persistence of LA over time (12 weeks or more between positive testing results) is a clinically important criterion for the antiphospholipid antibody syndrome diagnosis.

### Clinical Reference

1. Proven A, Bartlett RP, Moder KG, et al. Clinical importance of positive test results for lupus anticoagulant and anticardiolipin antibodies. *Mayo Clin Proc.* 2004;79(4):467-475
2. Gastineau DA, Kazmier FJ, Nichols WL, Bowie EJ. Lupus anticoagulant: an analysis of the clinical and laboratory features of 219 cases. *Am J Hematol.* 1985;19(3):265-275
3. Brandt JT, Triplett DA, Alving B, Sharrer I. Criteria for the diagnosis of lupus anticoagulant: an update. On behalf of the Subcommittee on Lupus Anticoagulant/Antiphospholipid Antibody of the Scientific and Standardisation Committee of the ISTH. *Thromb Haemost.* 1995;74(4):1185-1190
4. Arnout J, Vermeylen J. Current status and implications of autoimmune antiphospholipid antibodies in relation to thrombotic disease. *J Thromb Haemost.* 2003;1(5):931-942
5. Pengo V, Tripodi A, Reber G, Rand JH, et al. Update of the guidelines for lupus anticoagulant detection. Subcommittee on Lupus Anticoagulant/Antiphospholipid Antibody of the Scientific and Standardisation Committee of the International Society on Thrombosis and Haemostasis. *J Thromb Haemost.* 2009;7:1737-1740. doi:10.1111/j.1538-7836.2009.03555.x
6. CLSI Laboratory Testing for Lupus Anticoagulant; Approved Guideline. CLSI document H60-A. Clinical and Laboratory Standards Institute; 2014

### Performance

#### Method Description

The dilute Russell's viper venom time (DRVVT) mix assay is performed on the Instrumentation Laboratory ACL TOP. Patient plasma is mixed in a 1:1 ratio with normal pooled plasma and incubated for a specified time, then combined with a DRVVT screening reagent containing Russell's viper venom, phospholipids, heparin neutralizing agents, calcium, buffers, stabilizers, and sodium azide to trigger the coagulation process. Subsequently, the time to clot formation is measured optically using a wavelength of 671 nm.

The patient DRVVT mix clotting time is normalized by dividing the patient result by the mean DRVVT screening clotting time of normal pooled plasma to yield a ratio.(Package insert: CRYOcheck LA Check, PrecisionBioLogic, Inc.; Insert I-021 Rev. 11)

**PDF Report**

No

**Day(s) Performed**

Monday through Friday

**Report Available**

2 to 5 days

**Specimen Retention Time**

7 days

**Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

**Fees & Codes****Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**Test Classification**

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

**CPT Code Information**

85613

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
DRV2	DRVVT Mix	75512-4

Result ID	Test Result Name	Result LOINC® Value
RVMR2	DRVVT Mix Ratio	75512-4