

Targeted Benzodiazepine Screen, Random, Urine

#### Overview

#### **Useful For**

Determining compliance or identifying illicit benzodiazepine drug use

This test is **not intended for** employment-related testing.

#### **Profile Information**

Test Id	Reporting Name	Available Separately	Always Performed
LPPB	List prescribed	No	Yes
	benzodiazepines		
TABSU	Targeted Benzodiazepine	No	Yes
	Screen, U		

#### **Special Instructions**

• Clinical Toxicology CPT Code Client Guidance

# **Highlights**

This test uses high-resolution accurate mass spectrometry to identify 27 different benzodiazepines and metabolites for situations when immunoassays are not adequate.

This test offers lower detection limits than previous Mayo Clinic Laboratories-offered screens.

This test has the ability to detect "spiked" samples.

#### **Method Name**

Liquid Chromatography Tandem Mass Spectrometry, High-Resolution Accurate Mass (LC-MS/MS HRAM)

# **NY State Available**

Yes

# Specimen

# **Specimen Type**

Urine

# **Additional Testing Requirements**

In most cases, no additional testing is needed after the qualitative targeted benzodiazepine test is performed if the



Targeted Benzodiazepine Screen, Random, Urine

parent drug or metabolites found are consistent with the patient's prescribed medications. However, if unexpected benzodiazepine parent or metabolites are found, confirmatory testing can be requested at an additional charge.

## **Specimen Required**

Supplies: Sarstedt Aliquot Tube 5 mL (T914)

Collection Container/Tube: Plastic urine container

Submission Container/Tube: Plastic, 5-mL tube

**Specimen Volume:** 1 mL **Collection Instructions:** 

1. Collect a random urine specimen.

2. No preservative

#### **Forms**

If not ordering electronically, complete, print, and send a <u>Therapeutics Test Request</u> (T831) with the specimen.

#### Specimen Minimum Volume

0.5 mL

#### Reject Due To

Gross	Reject
hemolysis	
Gross icterus	Reject

#### **Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	14 days	
	Ambient	72 hours	
	Frozen	28 days	

## Clinical & Interpretive

#### **Clinical Information**

Benzodiazepines represent a large family of medications used to treat a wide range of disorders from anxiety to seizures, and they are also used in pain management. With a high risk for abuse and diversion, professional practice guidelines recommend compliance monitoring for these medications using urine drug tests. However, traditional benzodiazepine immunoassays suffer from a lack of cross-reactivity with all the benzodiazepines, so many compliant patients taking either clonazepam (Klonopin) or lorazepam (Ativan) may screen negative by immunoassay but are positive when confirmatory testing is done. The new targeted benzodiazepine screening test provides a more sensitive and specific test to check for compliance to all the commonly prescribed benzodiazepines and looks for both the parent drug and its metabolites in the urine.



Targeted Benzodiazepine Screen, Random, Urine

#### **Reference Values**

Not detected (Positive results are reported with qualitative "Present" results)

Cutoff concentrations: Alprazolam: 10 ng/mL

Alpha-hydroxyalprazolam: 10 ng/mL

Alpha-hydroxyalprazolam glucuronide: 50 ng/mL

Chlordiazepoxide: 10 ng/mL

Clobazam: 10 ng/mL

N-Desmethylclobazam: 200 ng/mL

Clonazepam: 10 ng/mL

7-Aminoclonazepam: 10 ng/mL

Diazepam: 10 ng/mL Nordiazepam: 10 ng/mL Flunitrazepam: 10 ng/mL

7-Aminoflunitrazepam: 10 ng/mL

Flurazepam: 10 ng/mL

2-Hydroxy ethyl flurazepam: 10 ng/mL

Lorazepam: 10 ng/mL

Lorazepam glucuronide: 50 ng/mL

Midazolam: 10 ng/mL

Alpha-hydroxy midazolam: 10 ng/mL

Oxazepam: 10 ng/mL

Oxazepam glucuronide: 50 ng/mL

Prazepam: 10 ng/mL Temazepam: 10 ng/mL

Temazepam glucuronide: 50 ng/mL

Triazolam: 10 ng/mL

Alpha-hydroxy triazolam: 10 ng/mL

Zolpidem: 10 ng/mL

Zolpidem phenyl-4-carboxylic acid: 10 ng/mL

#### Interpretation

If a benzodiazepine or its corresponding metabolites is identified (present), it indicates that the patient has used the respective benzodiazepine in the recent past. The absence of expected benzodiazepines or their metabolites may indicate noncompliance, inappropriate timing of specimen collection relative to drug administration, poor drug absorption, diluted or adulterated urine, or limitations of testing. The concentration of the drug must be greater than or equal to the cutoff to be reported as present. If a specific drug concentration is required, the laboratory must be contacted within 2 weeks of specimen collection/testing to request quantification by a second analytical technique at an additional charge.

#### Cautions

No significant cautionary statements

#### **Clinical Reference**



Targeted Benzodiazepine Screen, Random, Urine

- 1. Jannetto PJ, Bratanow NC, Clark WA, et al. Executive summary: American Association of Clinical Chemistry Laboratory Medicine Practice Guideline-Using clinical laboratory tests to monitor drug therapy in pain management patients. J Appl Lab Med. 2018;2(4):489-526
- 2. Langman LJ, Bechtel LK, Holstege CP. Clinical toxicology. In: Rifai N, Chiu RWK, Young I, Burnham CAD, Wittwer CT, eds. Tietz Textbook of Laboratory Medicine. 7th ed. Elsevier; 2023:chap 43

#### **Performance**

#### **Method Description**

The urine sample is diluted with internal standard and then analyzed by liquid chromatography tandem mass spectrometry using a high-resolution accurate mass orbitrap detector. (Unpublished Mayo method)

#### **PDF Report**

No

## Day(s) Performed

Monday through Sunday

#### Report Available

3 to 4 days

#### **Specimen Retention Time**

14 days

#### **Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Superior Drive

## **Fees & Codes**

#### **Fees**

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

#### **Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

#### **CPT Code Information**

G0480

80347 (if appropriate for select payers)

<u>Clinical Toxicology CPT Code Client Guidance</u>



Targeted Benzodiazepine Screen, Random, Urine

# **LOINC®** Information

Test ID	Test Order Name	Order LOINC® Value
TBSU	Targeted Benzodiazepine Screen, U	94117-9

Result ID	Test Result Name	Result LOINC® Value
604871	Alprazolam	94116-1
604867	Alpha-Hydroxyalprazolam	19325-0
604891	Alpha-Hydroxyalprazolam	94115-3
	Glucuronide	
604872	Chlordiazepoxide	19385-4
604889	Clobazam	94114-6
604890	N-Desmethylclobazam	94113-8
604873	Clonazepam	19399-5
604267	7-aminoclonazepam	94112-0
604874	Diazepam	19443-1
604880	Nordiazepam	19624-6
604875	Flunitrazepam	19466-2
604866	7-aminoflunitrazepam	94111-2
604876	Flurazepam	19474-6
604868	2-Hydroxy Ethyl Flurazepam	94110-4
604877	Lorazepam	19520-6
604878	Lorazepam Glucuronide	94109-6
604879	Midazolam	19585-9
604869	Alpha-Hydroxy Midazolam	94108-8
604881	Oxazepam	19638-6
604882	Oxazepam Glucuronide	94107-0
604883	Prazepam	19678-2
604884	Temazepam	19698-0
604885	Temazepam Glucuronide	94106-2
604886	Triazolam	19714-5
604870	Alpha-Hydroxy Triazolam	94105-4
604887	Zolpidem	94104-7
604888	Zolpidem Phenyl-4-Carboxylic acid	94103-9
604949	Benzodiazepine Interpretation	69050-3
LPPB	List prescribed benzodiazepines	29305-0