

Overview

Useful For

Investigation of possible diagnosis of Niemann-Pick disease types A and B

This test is **not recommended** for carrier detection because of the wide range of enzymatic activities observed in carriers and noncarriers.

Genetics Test Information

This test provides diagnostic testing for patients with decreased acid sphingomyelinase activity on newborn screen or with clinical signs and symptoms suspicious for Niemann-Pick type A or B.

Testing Algorithm

For more information see [Newborn Screen Follow-up for Acid Sphingomyelinase Deficiency](#)

If the patient has abnormal newborn screening results for Niemann- Pick disease, refer to the appropriate American College of Medical Genetics and Genomics Newborn Screening ACT Sheet.(1)

Special Instructions

- [Informed Consent for Genetic Testing](#)
- [Biochemical Genetics Patient Information](#)
- [Informed Consent for Genetic Testing \(Spanish\)](#)
- [Newborn Screen Follow-up for Acid Sphingomyelinase Deficiency](#)

Method Name

Flow Injection Analysis-Tandem Mass Spectrometry (FIA-MS/MS)

NY State Available

Yes

Specimen

Specimen Type

Whole Blood ACD

Shipping Instructions

For optimal isolation of leukocytes, it is recommended the specimen arrive refrigerated within 6 days of collection to be stabilized. Collect specimen Monday through Thursday only and not the day before a holiday. Specimen should be collected and packaged as close to shipping time as possible.

Specimen Required

Container/Tube:

**Preferred:** Yellow top (ACD solution B)  
**Acceptable:** Yellow top (ACD solution A) or lavender top (EDTA)  
**Specimen Volume:** 6 mL  
**Collection Instructions:** Send specimen in original tube. **Do not aliquot.**

Forms

1. **New York Clients-Informed consent is required.** Document on the request form or electronic order that a copy is on file. The following documents are available:
- [Informed Consent for Genetic Testing](#) (T576)
  - [Informed Consent for Genetic Testing-Spanish](#) (T826)
2. [Biochemical Genetics Patient Information](#) (T602)
3. If not ordering electronically, complete, print, and send a [Biochemical Genetics Test Request](#) (T798) with the specimen.

**Specimen Minimum Volume**  
2 mL

Reject Due To

Gross hemolysis	Reject
-----------------	--------

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Whole Blood ACD	Refrigerated (preferred)	6 days	
	Ambient	6 days	

Clinical & Interpretive

Clinical Information

Niemann-Pick disease (NPD) types A (NPA) and B (NPB) are autosomal recessive lysosomal storage disorders affecting metabolism of specific lipids within cells. NPA and NPB are caused by a deficiency of sphingomyelinase, which results in extensive storage of sphingomyelin and cholesterol in the liver, spleen, lungs, and, to a lesser degree, brain. NPA disease is more severe than NPB and is characterized by early onset with feeding problems, dystrophy, persistent jaundice, development of hepatosplenomegaly, neurological deterioration, deafness, and blindness, leading to death by age 3. NPB disease is limited to visceral symptoms with survival into adulthood. Some patients have been described with intermediary phenotypes. Large lipid-laden foam cells are characteristic of the disease. Approximately 50% of cases have cherry-red spots in the macula.

The combined prevalence of NPA and NPB is estimated to be 1 in 250,000. NPA and NPB are inherited in an autosomal recessive manner and are caused by variants in the *SMPD1* gene. Although there is a higher frequency of type A among the Ashkenazi Jewish population, both types are panethnic. Individuals with NPD types A and B typically have elevations of lyso-sphingomyelin and lyso-sphingomyelin 509 combined with elevations of the oxysterols cholestane-3 beta, 5

alpha, 6 beta-triol (COT) and 7-ketocholesterol (7-KC). (OXNP / Oxysterols, Plasma; OXYWB / Oxysterols, Blood; OXYBS / Oxysterols, Blood Spot). Molecular genetic testing for NPA and NPB disease is also available (CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies; specificity *SMPD1* Gene List ID: IEMCP-W6S9XD).

**Reference Values**

> or =0.32 nmol/hour/mg protein

An interpretative report will be provided.

**Interpretation**

Values below the reference range are consistent with a diagnosis for Niemann-Pick types A and B.

When abnormal results are detected, a detailed interpretation is given, including an overview of the results and of their significance, a correlation to available clinical information, elements of differential diagnosis, recommendations for additional biochemical testing, and in vitro, confirmatory studies (enzyme assay, molecular analysis), name and phone number of key contacts who may provide these studies, and a phone number to reach one of the laboratory directors in case the referring physician has additional questions.

**Cautions**

This test can give false-positive acid sphingomyelinase results. OXYBS / Oxysterols, Blood Spot may be ordered as a confirmatory test.

Additional biochemical or molecular testing is recommended to confirm a diagnosis if an enzyme deficiency is detected by this screening test.

**Clinical Reference**

1. Newborn Screening ACT Sheet [Decreased acid sphingomyelinase] Acid Sphingomyelinase Deficiency (ASMD). American College of Medical Genetics and Genomics; 2022. Revised May 2022. Accessed June 10, 2024. Available at [www.acmg.net/PDFLibrary/Niemann-Pick.pdf](http://www.acmg.net/PDFLibrary/Niemann-Pick.pdf)
2. Elliott S, Buroker N, Cournoyer JJ, et al: Pilot study of newborn screening for six lysosomal storage diseases using Tandem Mass Spectrometry. *Mol Genet Metab*. 2016 Aug;118(4):304-309
3. Matern D, Gavrilov D, Oglesbee D, Raymond K, Rinaldo P, Tortorelli S: Newborn screening for lysosomal storage disorders. *Semin Perinatol*. 2015 Apr;39(3):206-216
4. Schuchman EH, Desnick RJ: Niemann-Pick disease types A and B: acid sphingomyelinase deficiencies. In: Valle D, Antonarakis S, Ballabio A, Beaudet A, Mitchell GA, eds. *The Online Metabolic Bases of Inherited Disease*. McGraw-Hill; 2019. Accessed May 26, 2021. Available at <https://ommbid.mhmedical.com/content.aspx?sectionid=225545671&bookid=2709>
5. Wasserstein MP, Schuchman EH: Acid sphingomyelinase deficiency. In: Adam MP, Ardinger HH, Pagon RA, et al, eds. *GeneReviews*. [Internet]. University of Washington, Seattle; 2006. Updated February 25, 2021. Accessed May 26, 2021. Available at [www.ncbi.nlm.nih.gov/books/NBK1370/](http://www.ncbi.nlm.nih.gov/books/NBK1370/)
6. Schuchman EH, Desnick RJ: Types A and B Niemann-Pick disease. *Mol Genet Metab*. 2017 Jan-Feb;120(1-2):27-33

**Performance****Method Description**

The specimens are incubated with a mix of substrate and internal standard for acid sphingomyelinase ,beta-glucocerebrosidase , acid alpha-glucosidase , alpha-galactosidase , galactocerebrosidase and alpha-L-iduronidase . The sample is then purified by liquid-liquid extraction. The extract is evaporated and reconstituted before analysis by tandem mass spectrometry.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Preanalytical processing: Monday through Saturday.

Assay performed: Monday, Wednesday

Report Available

5 to 10 days

Specimen Retention Time

WBC homogenate: 1 month

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

- Fees
- Authorized users can sign in to [Test Prices](#) for detailed fee information.
  - Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
  - Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

82657

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
ASMW	Acid Sphingomyelinase, Leukocytes	24101-8

Result ID	Test Result Name	Result LOINC® Value
606264	Acid Sphingomyelinase, Leukocytes	24101-8
606265	Interpretation	59462-2
606266	Reviewed By	18771-6