

Test Definition: 7INHT

Factor VII Inhibitor Profile, Technical Interpretation

| _ | | | | |
|---|----|----|-----|---|
| n | ve | rv | Ie. | w |

Useful For

Technical interpretation of inhibitor to coagulation factor VII testing

Method Name

Only orderable as part of a profile. For more information see 7INHE / Factor VII Inhibitor Evaluation, Plasma.

Technical Interpretation

NY State Available

Yes

Specimen

Specimen Type

Plasma Na Cit

Specimen Required

Only orderable as part of a profile. For more information see 7INHE / Factor VII Inhibitor Evaluation, Plasma.

Reject Due To

| Gross | Reject |
|---------------|--------|
| hemolysis | |
| Gross lipemia | Reject |
| Gross icterus | Reject |

Specimen Stability Information

| Specimen Type | Temperature | Time | Special Container |
|---------------|-------------|---------|-------------------|
| Plasma Na Cit | Frozen | 14 days | |

Clinical & Interpretive

Clinical Information

Coagulation factor inhibitors arise in patients who are congenitally deficient in a specific factor in response to factor replacement therapy or can occur spontaneously without known cause or in response to a variety of medical conditions,



Test Definition: 7INHT

Factor VII Inhibitor Profile, Technical Interpretation

including the postpartum state, immunologic disorders, certain antibiotic therapies, some malignancies, and in the older population.

Inhibitors of factor VIII coagulant activity are the most commonly occurring of the specific factor inhibitors.

Reference Values

Only orderable as part of a profile. For more information see 7INHE / Factor VII Inhibitor Evaluation, Plasma.

An interpretive report will be provided.

Interpretation

When testing is complete, if factor activity results fall within clinically normal ranges, an interpretive comment will be provided noting that inhibitor testing was not indicated and, therefore, not performed. If factor activity indicates the performance of inhibitor screen testing, an interpretive comment will be provided noting the presence or absence of a factor VII inhibitor.

Cautions

Occasionally, a potent lupus-like anticoagulant may cause false-positive testing for a specific factor inhibitor (eg, factor VIII or IX).

Clinical Reference

- 1. Hoffman R, Benz Jr EJ, Silberstein LE, et al, eds. Hematology: Basic Principles and Practice. 7th ed. Elsevier; 2018
- 2. Kasper CK: Treatment of factor VIII inhibitors. Prog Hemost Thromb. 1989;9:57-86
- 3. Kottke-Marchant K, ed. Laboratory Hematology Practice. Wiley Blackwell Publishing; 2012

Performance

Method Description

If the factor VII activity level is normal or elevated, a computer-generated interpretive comment indicating inhibitor testing was not indicated and was not performed will be provided.

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

3 to 5 days

Specimen Retention Time

7 days

Performing Laboratory Location



Test Definition: 7INHT

Factor VII Inhibitor Profile, Technical Interpretation

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

Test Classification

Not Applicable

CPT Code Information

85390

LOINC® Information

| Test ID | Test Order Name | Order LOINC® Value |
|---------|--------------------------------|--------------------|
| 7INHT | FVII Inhib Profile Tech Interp | 69049-5 |
| | | |

| Result ID | Test Result Name | Result LOINC® Value |
|-----------|--------------------------------|---------------------|
| 7INHT | FVII Inhib Profile Tech Interp | 69049-5 |