

Overview

Useful For

Follow up for abnormal biochemical or electron microscopy results suspicious for neuronal ceroid lipofuscinoses (NCL)

Establishing a molecular diagnosis for patients with NCL

Identifying variations within genes known to be associated with NCL, allowing for predictive testing of at-risk family members

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
CULFB	Fibroblast Culture for Genetic Test	Yes	No

Genetics Test Information

This test utilizes next-generation sequencing to detect single nucleotide and copy number variants in 16 genes associated with neuronal ceroid lipofuscinosis (NCL/Batten Disease): *ATP13A2, CLN3, CLN5, CLN6, CLN8, CTSD, CTSF, CTSK, DNAJC5, GRN, KCTD7, MFSD8, PANK2, PPT1, SGSH, TPP1*. See [Targeted Genes and Methodology Details for Neuronal Ceroid Lipofuscinosis \(Batten Disease\) Gene Panel](#) in Special Instructions and Method Description for additional details.

Identification of a pathogenic variant may assist with diagnosis, prognosis, clinical management, familial screening, and genetic counseling for NCL (Batten disease).

Additional first tier testing may be considered/recommended. For more information see Ordering Guidance.

Testing Algorithm

For skin biopsy or cultured fibroblast specimens, fibroblast culture testing will be performed at an additional charge. If viable cells are not obtained, the client will be notified.

Special Instructions

- [Molecular Genetics: Biochemical Disorders Patient Information](#)
- [Informed Consent for Genetic Testing](#)
- [Blood Spot Collection Card-Spanish Instructions](#)
- [Blood Spot Collection Card-Chinese Instructions](#)
- [Informed Consent for Genetic Testing \(Spanish\)](#)
- [Blood Spot Collection Instructions](#)
- [Targeted Genes and Methodology Details for Neuronal Ceroid Lipofuscinosis \(Batten Disease\) Gene Panel](#)

Method Name

Sequence Capture and Targeted Next-Generation Sequencing followed by Polymerase Chain Reaction (PCR) and Sanger Sequencing

**NY State Available**

Yes

**Specimen**

**Specimen Type**

Varies

**Ordering Guidance**

First-tier biochemical testing is available for the 2 most common types of enzyme deficiency associated with neuronal ceroid lipofuscinosis; order TPPTL / Tripeptidyl Peptidase 1 and Palmitoyl-Protein Thioesterase 1, Leukocytes.

Customization of this panel and single gene analysis for any gene present on this panel is available. For more information see CGPH/ Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies.

**Shipping Instructions**

Specimen preferred to arrive within 96 hours of collection.

**Specimen Required**

**Patient Preparation:** A previous bone marrow transplant from an allogenic donor will interfere with testing. Call 800-533-1710 for instructions for testing patients who have received a bone marrow transplant.

**Submit only 1 of the following specimens:**

**Specimen Type:** Whole blood

**Container/Tube:**

**Preferred:** Lavender top (EDTA) or yellow top (ACD)

**Acceptable:** Any anticoagulant

**Specimen Volume:** 3 mL

**Collection Instructions:**

1. Invert several times to mix blood.
2. Send whole blood specimen in original tube. **Do not aliquot.**

**Specimen Stability Information:** Ambient (preferred) 4 days/Refrigerated 14 days

**Specimen Type:** Skin biopsy

**Supplies:** Fibroblast Biopsy Transport Media (T115)

**Container/Tube:** Sterile container with any standard cell culture media (eg, minimal essential media, RPMI 1640). The solution should be supplemented with 1% penicillin and streptomycin.

**Specimen Volume:** 4-mm punch

---

**Specimen Stability Information:** Refrigerated (preferred)/Ambient

**Additional Information:** A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks is required to culture fibroblasts before genetic testing can occur.

**Specimen Type:** Cultured fibroblast

**Container/Tube:** T-25 flask

**Specimen Volume:** 2 Flasks

**Collection Instructions:** Submit confluent cultured fibroblast cells from a skin biopsy from another laboratory. Cultured cells from a prenatal specimen will not be accepted.

**Specimen Stability Information:** Ambient (preferred)/Refrigerated (<24 hours)

**Additional Information:** A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks is required to culture fibroblasts before genetic testing can occur.

**Specimen Type:** Blood spot

**Supplies:** Card-Blood Spot Collection (Filter Paper) (T493)

**Container/Tube:**

**Preferred:** Collection card (Whatman Protein Saver 903 Paper)

**Acceptable:** PerkinElmer 226 (formerly Ahlstrom 226) filter paper or blood spot collection card

**Specimen Volume:** 5 Blood spots

**Collection Instructions:**

1. An alternative blood collection option for a patient older than 1 year is a fingerstick. For detailed instructions, see [How to Collect Dried Blood Spot Samples](#).
2. Let blood dry on the filter paper at ambient temperature in a horizontal position for a minimum of 3 hours.
3. Do not expose specimen to heat or direct sunlight.
4. Do not stack wet specimens.
5. Keep specimen dry

**Specimen Stability Information:** Ambient (preferred)/Refrigerated

**Additional Information:**

1. Due to lower concentration of DNA yielded from blood spot, it is possible that additional specimen may be required to complete testing.
2. For collection instructions, see [Blood Spot Collection Instructions](#)
3. For collection instructions in Spanish, see [Blood Spot Collection Card-Spanish Instructions](#) (T777)
4. For collection instructions in Chinese, see [Blood Spot Collection Card-Chinese Instructions](#) (T800)

**Specimen Type:** Saliva

**Patient Preparation:** Patient should not eat, drink, smoke, or chew gum 30 minutes prior to collection.

**Supplies:** Saliva Swab Collection Kit (T786)

**Specimen Volume:** 1 Swab

**Collection Instructions:** Collect and send specimen per kit instructions.

**Specimen Stability Information:** Ambient 30 days

**Additional Information:** Due to lower concentration of DNA yielded from saliva, it is possible that additional specimen may be required to complete testing.

## Forms

1. **New York Clients-Informed consent is required.** Document on the request form or electronic order that a copy is on

file. The following documents are available:

- [Informed Consent for Genetic Testing](#) (T576)
- [Informed Consent for Genetic Testing \(Spanish\)](#) (T826)
- 2. [Molecular Genetics: Biochemical Disorders Patient Information](#) (T527)
- 3. If not ordering electronically, complete, print, and send a [Biochemical Genetics Test Request](#) (T798) with the specimen.

Specimen Minimum Volume

See Specimen Required

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Varies	Varies		

Clinical & Interpretive

Clinical Information

Neuronal ceroid lipofuscinoses (NCL) are a subset of lysosomal storage diseases that involve defective cellular processing of lipids. NCL are clinically characterized by epilepsy, intellectual and motor decline, and blindness. Electron microscopy typically shows a characteristic accumulation of granular osmophilic deposits (GROD), curvilinear profiles (CVB), or fingerprint profiles (FP). Enzymatic testing may show deficiency in palmitoyl-protein thioesterase 1 (PPT1), tripeptidyl-peptidase 1 (TPP1), or cathepsin D (CTSD). Currently there are at least 14 genetically distinct forms.

Age of onset and clinical features can be variable, from congenital to adult onset. NCL is typically inherited in an autosomal recessive manner, although one adult onset form (ANCL; *DNAJC5* gene) has been shown to be autosomal dominant.

Reference Values

An interpretive report will be provided.

Interpretation

All detected alterations are evaluated according to American College of Medical Genetics and Genomics (ACMG) recommendations.(1) Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.

Cautions

Clinical Correlations:  
Test results should be interpreted in the context of clinical findings, family history, and other laboratory data. Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

---

If testing was performed because of a clinically significant family history, it is often useful to first test an affected family member. Detection of at least one reportable variant in an affected family member would allow for more informative testing of at-risk individuals.

To discuss the availability of additional testing options or for assistance in the interpretation of these results, contact the Mayo Clinic Laboratory genetic counselors at 800-533-1710.

**Technical Limitations:**

Next-generation sequencing may not detect all types of genomic variants. In rare cases, false-negative or false-positive results may occur. The depth of coverage may be variable for some target regions; assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be considered.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. Confirmation of select reportable variants will be performed by alternate methodologies based on internal laboratory criteria.

This test is validated to detect 95% of deletions up to 75 base pairs (bp) and insertions up to 47 bp. Insertions/deletions (indels) of 40 or more bp, including mobile element insertions, may be less reliably detected than smaller indels.

**Deletion/Duplication Analysis:**

This analysis targets single and multi-exon deletions/duplications; however, in some instances single exon resolution cannot be achieved due to isolated reduction in sequence coverage or inherent genomic complexity. Balanced structural rearrangements (such as translocations and inversions) may not be detected.

This test is not designed to detect low levels of mosaicism or to differentiate between somatic and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

Genes may be added or removed based on updated clinical relevance. Refer to the [Targeted Genes and Methodology Details for Neuronal Ceroid Lipofuscinosis \(Batten Disease\) Gene Panel](#) in Special Instructions for the most up to date list of genes included in this test. For detailed information regarding gene specific performance and technical limitations, see Method Description or contact a laboratory genetic counselor.

If the patient has had an allogeneic hematopoietic stem cell transplant or a recent heterologous blood transfusion, results may be inaccurate due to the presence of donor DNA. Call Mayo Clinic Laboratories for instructions for testing patients who have received a bone marrow transplant.

**Reclassification of Variants:**

At this time, it is not standard practice for the laboratory to systematically review previously classified variants on a regular basis. The laboratory encourages health care providers to contact the laboratory at any time to learn how the classification of a particular variant may have changed over time.

**Variant Evaluation:**

Evaluation and categorization of variants is performed using published American College of Medical Genetics and Genomics and the Association for Molecular Pathology recommendations as a guideline. Other gene-specific guidelines may also be considered. Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance. Variants classified as benign or likely benign are not reported.

Multiple in silico evaluation tools may be used to assist in the interpretation of these results. The accuracy of predictions made by in silico evaluation tools is highly dependent upon the data available for a given gene, and periodic updates to these tools may cause predictions to change over time. Results from in silico evaluation tools should be interpreted with caution and professional clinical judgement.

**Clinical Reference**

1. Richards S, Aziz N, Bale S, et al: Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. Genet Med. 2015 May;17(5):405-424
2. Mole SE, Cotman SL: Genetics of the neuronal ceroid lipofuscinoses (Batten disease). Biochim Biophys Acta. 2015;1852(10PtB):2237-2241
3. Cooper JD, Tarczyk MA, Nelvagal HR: Towards a new understanding of NCL pathogenesis. Biochim Biophys Acta. 2015;1852(10PtB):2256-2261
4. Williams RE, Mole SE: New nomenclature and classification scheme for the neuronal ceroid lipofuscinoses. Neurology. 2012;79(2):183-191
5. Mole SE, Williams RE: Neuronal ceroid-lipofuscinoses. In: Adam MP, Ardinger HH, Pagon RA, et al, eds. GeneReviews [Internet]. University of Washington. , Seattle; 2001. Updated August 1, 2013. Accessed October 28, 2020. Available at [www.ncbi.nlm.nih.gov/books/NBK1428/](http://www.ncbi.nlm.nih.gov/books/NBK1428/)

**Performance****Method Description**

Next-generation sequencing (NGS) and/or Sanger sequencing is performed to test for the presence of variants in coding regions and intron/exon boundaries of the genes analyzed, as well as some other regions that have known pathogenic variants. The human genome reference GRCh37/hg19 build was used for sequence read alignment. At least 99% of the bases are covered at a read depth over 30X. Sensitivity is estimated at above 99% for single nucleotide variants, above 94% for insertions/deletions (indels) less than 40 base pairs (bp), above 95% for deletions up to 75 bp and insertions up to 47 bp. NGS and/or a polymerase chain reaction (PCR)-based quantitative method is performed to test for the presence of deletions and duplications in the genes analyzed. See [Targeted Genes and Methodology Details for Neuronal Ceroid Lipofuscinosis \(Batten Disease\) Gene Panel](#) in Special Instructions for details regarding the targeted genes analyzed.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis

as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. See [Targeted Genes and Methodology Details for Neuronal Ceroid Lipofuscinosis \(Batten Disease\) Gene Panel](#) in Special Instructions for details regarding the specific gene regions not routinely covered.(Unpublished Mayo method)

Genes analyzed: *ATP13A2, CLN3, CLN5, CLN6, CLN8, CTSD, CTSF, CTSK, DNAJC5, GRN, KCTD7, MFSD8, PANK2, PPT1, SGSH, TPP1*

PDF Report  
Supplemental

Day(s) Performed  
Varies

Report Available  
28 to 42 days

Specimen Retention Time  
Whole Blood: 2 weeks (if available); Extracted DNA: 3 months; Blood spots/Saliva; 1 month

Performing Laboratory Location  
Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

- Fees
- Authorized users can sign in to [Test Prices](#) for detailed fee information.
  - Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
  - Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification  
This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information  
81443  
88233-Tissue culture, skin, solid tissue biopsy (if appropriate)  
88240-Cryopreservation (if appropriate)

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
---------	-----------------	--------------------

Test Definition: NCLGP

Neuronal Ceroid Lipofuscinosis (Batten Disease) Gene Panel, Varies

NCLGP	NCL (Batten Disease) Gene Panel	105349-5
-------	---------------------------------	----------

Result ID	Test Result Name	Result LOINC® Value
608572	Test Description	62364-5
608573	Specimen	31208-2
608574	Source	31208-2
608575	Result Summary	50397-9
608576	Result	82939-0
608577	Interpretation	69047-9
608578	Resources	99622-3
608579	Additional Information	48767-8
608580	Method	85069-3
608581	Genes Analyzed	48018-6
608582	Disclaimer	62364-5
608583	Released By	18771-6