

## Overview

### Useful For

Follow up for abnormal biochemical results suggestive of a urea cycle disorder (UCD)

Establishing a molecular diagnosis for patients with a UCD

Identifying variants within genes known to be associated with UCD, allowing for predictive testing of at-risk family members

### Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
CULFB	Fibroblast Culture for Genetic Test	Yes	No

### Genetics Test Information

This test utilizes next-generation sequencing to detect single nucleotide and copy number variants in 16 genes associated with urea cycle disorders (UCD). This test includes next-generation sequencing to test for variants in the genes indicated: *ALDH18A1*, *ARG1*, *ARG2*, *ASL*, *ASS1*, *CA5A*, *CPS1*, *GLUD1*, *GLUL*, *NAGS*, *OAT*, *OTC*, *SLC25A13*, *SLC25A15*, *SLC7A7*, and *UMPS*. See [Targeted Genes and Methodology Details for Urea Cycle Disorders Gene Panel](#) and Method Description for additional details.

Identification of a disease-causing variant may assist with diagnosis, prognosis, clinical management, familial screening, and genetic counseling for UCD.

### Testing Algorithm

For skin biopsy and cultured fibroblast specimens, fibroblast culture will be added at an additional charge. If viable cells are not obtained, the client will be notified.

### Special Instructions

- [Molecular Genetics: Biochemical Disorders Patient Information](#)
- [Informed Consent for Genetic Testing](#)
- [Blood Spot Collection Card-Spanish Instructions](#)
- [Blood Spot Collection Card-Chinese Instructions](#)
- [Informed Consent for Genetic Testing \(Spanish\)](#)
- [Blood Spot Collection Instructions](#)
- [Targeted Genes and Methodology Details for Urea Cycle Disorders Gene Panel](#)

### Method Name

Sequence Capture and Targeted Next-Generation Sequencing (NGS) followed by Polymerase Chain Reaction (PCR) and Sanger Sequencing

### NY State Available

---

Yes

## Specimen

### Specimen Type

Varies

### Ordering Guidance

The recommended first-tier biochemical tests to screen for urea cycle disorders are a combination of quantitative plasma amino acids and urinary orotic acid. Order AAQP / Amino Acids, Quantitative, Plasma and OROT / Orotic Acid, Random, Urine.

Customization of this panel and single gene analysis for any gene present on this panel is available. For more information see CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies.

Targeted testing for familial variants (also called site-specific or known mutations testing) is available for the genes on this panel. See FMTT / Familial Variant, Targeted Testing, Varies. To obtain more information about this testing option, call 800-533-1710.

### Shipping Instructions

Specimen preferred to arrive within 96 hours of collection.

### Specimen Required

**Patient Preparation:** A previous bone marrow transplant from an allogenic donor will interfere with testing. For instructions for testing patients who have received a bone marrow transplant, call 800-533-1710.

### Submit only 1 of the following specimens:

**Specimen Type:** Whole blood

**Container/Tube:** Lavender top (EDTA) or yellow top (ACD)

**Specimen Volume:** 3 mL

**Collection Instructions:**

1. Invert several times to mix blood.
2. Send whole blood specimen in original tube. **Do not aliquot.**

**Specimen Stability Information:** Ambient (preferred) 4 days/Refrigerated 14 days

**Specimen Type:** Skin biopsy

**Supplies:** Fibroblast Biopsy Transport Media (T115)

**Container/Tube:** Sterile container with any standard cell culture media (eg, minimal essential media, RPMI 1640). The solution should be supplemented with 1% penicillin and streptomycin (Eagle's minimum essential medium with 1% penicillin and streptomycin).

**Specimen Volume:** 4-mm punch

**Specimen Stability Information:** Refrigerated (preferred)/Ambient

**Additional Information:** A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks is required to culture fibroblasts before genetic testing can occur.

**Specimen Type:** Cultured fibroblast

**Container/Tube:** T-25 flask

**Specimen Volume:** 2 Flasks

**Collection Instructions:** Submit confluent cultured fibroblast cells from a skin biopsy from another laboratory. Cultured cells from a prenatal specimen will not be accepted.

**Specimen Stability Information:** Ambient (preferred)/Refrigerated (<24 hours)

**Additional Information:** A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks is required to culture fibroblasts before genetic testing can occur.

**Specimen Type:** Blood spot

**Supplies:** Card-Blood Spot Collection (Filter Paper) (T493)

**Container/Tube:**

**Preferred:** Collection card (Whatman Protein Saver 903 Paper)

**Acceptable:** PerkinElmer 226 (formerly Ahlstrom 226) filter paper, or blood spot collection card

**Specimen Volume:** 5 Blood spots

**Collection Instructions:**

1. An alternative blood collection option for a patient older than 1 year is a fingerstick. For detailed instructions, see [How to Collect Dried Blood Spot Samples](#).
2. Let blood dry on the filter paper at ambient temperature in a horizontal position for a minimum of 3 hours.
3. Do not expose specimen to heat or direct sunlight.
4. Do not stack wet specimens.
5. Keep specimen dry.

**Specimen Stability Information:** Ambient (preferred)/Refrigerated

**Additional Information:**

1. Due to lower concentration of DNA yielded from blood spot, it is possible that additional specimen may be required to complete testing.
2. For collection instructions, see [Blood Spot Collection Instructions](#).
3. For collection instructions in Spanish, see [Blood Spot Collection Card-Spanish Instructions](#) (T777).
4. For collection instructions in Chinese, see [Blood Spot Collection Card-Chinese Instructions](#) (T800).

**Specimen Type:** Saliva

**Patient Preparation:** Patient should not eat, drink, smoke, or chew gum 30 minutes prior to collection.

**Supplies:**

DNA Saliva Kit High Yield (T1007)

Saliva Swab Collection Kit (T786)

**Container/Tube:**

**Preferred:** High-yield DNA saliva kit

**Acceptable:** Saliva swab

**Specimen Volume:** 1 Tube if using T1007 or 2 swabs if using T786

**Collection Instructions:** Collect and send specimen per kit instructions.

**Specimen Stability Information:** Ambient (preferred) 30 days/Refrigerated 30 days

**Additional Information:** Saliva specimens are acceptable but not recommended. Due to lower quantity/quality of DNA

yielded from saliva, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.

**Forms**

- New York Clients-Informed consent is required.** Document on the request form or electronic order that a copy is on file. The following documents are available:  
[Informed Consent for Genetic Testing \(T576\)](#)  
[Informed Consent for Genetic Testing \(Spanish\) \(T826\)](#)
- [Molecular Genetics: Biochemical Disorders Patient Information \(T527\)](#)
- If not ordering electronically, complete, print, and send a [Biochemical Genetics Test Request \(T798\)](#) with the specimen.

**Specimen Minimum Volume**

See Specimen Required

**Reject Due To**

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

**Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Varies	Varies		

**Clinical & Interpretive****Clinical Information**

Urea cycle disorders (UCD) are a group of inherited disorders of nitrogen detoxification that result when any of the enzymes in the urea cycle have reduced or absent activity. These disorders include carbamoylphosphate synthetase I deficiency, ornithine transcarbamylase (OTC) deficiency, argininosuccinic acid synthetase deficiency (citrullinemia), argininosuccinic acid lyase deficiency (argininosuccinic aciduria), arginase deficiency, the cofactor producer, N-acetyl glutamate synthetase (NAGS) deficiency, and two amino acid transporters, ornithine translocase deficiency (hyperornithinemia, hyperammonemia, homocitrullinuria syndrome) and citrin deficiency. The role of the urea cycle is to metabolize and clear waste nitrogen, and defects in any of the steps of the pathway can result in an accumulation of toxic ammonia in the nervous system. The urea cycle is also responsible for endogenous production of the amino acids citrulline, ornithine, and arginine.

Infants with a complete urea cycle enzyme deficiency typically appear normal at birth, but they present in the neonatal period with lethargy, seizures, hyper- or hypoventilation, and ultimately coma or death, as a result of elevated ammonia levels. Individuals with partial enzyme deficiency may present later in life, typically following an acute illness or other catabolic stressor. Symptoms may be less severe and may appear as episodes of psychosis, lethargy, cyclical vomiting, and behavioral abnormalities. Patients with impaired ornithine metabolism due to ornithine aminotransferase deficiency may present with childhood onset myopia progressing to vision loss in the 4th to 6th decades of life. Patients may or may not have accompanying hyperammonemia but display marked elevations in plasma ornithine.

---

All the UCD are inherited as autosomal recessive disorders, with the exception of OTC deficiency, which is X-linked. UCD may be suspected with elevated ammonia, normal anion gap, and a normal glucose.

This comprehensive gene panel is a helpful tool to establish a diagnosis for patients with suggestive clinical and biochemical features given the broad clinical spectrum and genetic heterogeneity of UCD. Molecular genetic testing can help to distinguish among the conditions and allows for diagnostic confirmation.

A combination of biochemical tests including quantitative plasma amino acids (AAQP / Amino Acids, Quantitative, Plasma) and urinary orotic acid (OROT / Orotic Acid, Random, Urine) are recommended as the first-tier test to assess patients for UCD.

Acute treatment for UCD consists of dialysis and administration of nitrogen scavenger drugs to reduce ammonia concentration. Chronic management typically involves restriction of dietary protein with essential amino acid supplementation. More recently, liver transplantation has been used with success in treating some patients.

## Reference Values

An interpretive report will be provided.

## Interpretation

All detected alterations are evaluated according to American College of Medical Genetics and Genomics recommendations.(1) Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.

## Cautions

### Clinical Correlations:

Test results should be interpreted in the context of clinical findings, family history, and other laboratory data. Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

If testing was performed because of a clinically significant family history, it is often useful to first test an affected family member. Detection of at least one reportable variant in an affected family member would allow for more informative testing of at-risk individuals.

To discuss the availability of additional testing options or for assistance in the interpretation of these results, contact the Mayo Clinic Laboratories genetic counselors at 800-533-1710.

### Technical Limitations:

Next-generation sequencing may not detect all types of genomic variants. In rare cases, false-negative or false-positive results may occur. The depth of coverage may be variable for some target regions; assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be considered.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. Confirmation of select reportable variants will be performed by alternate methodologies based on internal laboratory criteria.

---

This test is validated to detect 95% of deletions up to 75 base pairs (bp) and insertions up to 47 bp. Deletions-insertions (delins) of 40 or more bp, including mobile element insertions, may be less reliably detected than smaller delins.

**Deletion/Duplication Analysis:**

This analysis targets single and multi-exon deletions/duplications; however, in some instances single exon resolution cannot be achieved due to isolated reduction in sequence coverage or inherent genomic complexity. Balanced structural rearrangements (such as translocations and inversions) may not be detected.

This test is not designed to detect low levels of mosaicism or to differentiate between somatic and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

Genes may be added or removed based on updated clinical relevance. For detailed information regarding gene-specific performance and technical limitations, see Method Description or contact a laboratory genetic counselor.

If the patient has had an allogeneic hematopoietic stem cell transplant or a recent blood transfusion, results may be inaccurate due to the presence of donor DNA. Call Mayo Clinic Laboratories for instructions for testing patients who have received a bone marrow transplant.

**Reclassification of Variants:**

Currently, it is not standard practice for the laboratory to systematically review previously classified variants on a regular basis. The laboratory encourages healthcare providers to contact the laboratory at any time to learn how the classification of a particular variant may have changed over time. Due to broadening genetic knowledge, it is possible that the laboratory may discover new information of relevance to the patient. Should that occur, the laboratory may issue an amended report.

**Variant Evaluation:**

Evaluation and categorization of variants are performed using published American College of Medical Genetics and Genomics and the Association for Molecular Pathology recommendations as a guideline.(1) Other gene-specific guidelines may also be considered. Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance. Variants classified as benign or likely benign are not reported.

Multiple in silico evaluation tools may be used to assist in the interpretation of these results. The accuracy of predictions made by in silico evaluation tools is highly dependent upon the data available for a given gene, and periodic updates to these tools may cause predictions to change over time. Results from in silico evaluation tools should be interpreted with caution and professional clinical judgment.

Rarely, incidental or secondary findings may implicate another predisposition or presence of active disease. These findings will be carefully reviewed to determine whether they will be reported.

**Clinical Reference**

1. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular

Pathology. Genet Med. 2015;17(5):405-424

2. Brusilow SW, Horwitz AL. Urea cycle enzymes. In: Valle D, Antonarakis S, Ballabio A, Beaudet A, Mitchell GA. eds. The Online Metabolic and Molecular Bases of Inherited Disease. McGraw-Hill Education; 2019. Accessed March 8, 2024. Available at <https://ommbid.mhmedical.com/content.aspx?sectionid=225084071&bookid=2709&Resultclick=2>

3. Haberle J, Burlina A, Chakrapani A, et al. Suggested guidelines for diagnosis and management of urea cycle disorders: First revision. J Inherit Metab Dis. 2019;1-39. doi:10.1002/jimd.12100

4. Valle D, Simell O. The hyperornithinemias. In: Valle D, Antonarakis S, Ballabio A, Beaudet A, Mitchell GA. eds. The Online Metabolic and Molecular Bases of Inherited Disease. McGraw-Hill Education; 2019. Accessed March 8, 2024. Available at <https://ommbid.mhmedical.com/content.aspx?sectionid=225083672&bookid=2709&Resultclick=2>

5. Foshci FG, Morelli MC, Savini S, et al. Urea cycle disorders: A case report of a successful liver transplant and a literature review. World J Gastroenterol. 2015;21(13):4063-4068

6. Ah Mew N, Simpson KL, Gropman AL, et al. Urea cycle disorders overview. In: Adam MP, Everman DB, Mirzaa GM, et al. eds. GeneReviews [Internet]. University of Washington, Seattle; 2003. Updated June 22, 2017. Accessed March 8, 2024. Available at [www.ncbi.nlm.nih.gov/books/NBK1217/](http://www.ncbi.nlm.nih.gov/books/NBK1217/)

## Performance

### Method Description

Next-generation sequencing (NGS) and/or Sanger sequencing are performed to test for the presence of variants in coding regions and intron/exon boundaries of the genes analyzed, as well as some other regions that have known disease-causing variants. The human genome reference GRCh37/hg19 build was used for sequence read alignment. At least 99% of the bases are covered at a read depth over 30X. Sensitivity is estimated to be over 99% for single nucleotide variants, over 94% for deletions-insertions (delins) less than 40 base pairs (bp), and over 95% for deletions up to 75 bp and insertions up to 47 bp. NGS and/or a polymerase chain reaction-based quantitative method is performed to test for the presence of deletions and duplications in the genes analyzed.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. See [Targeted Genes and Methodology Details for Urea Cycle Disorders Gene Panel](#) for details regarding the targeted genes analyzed and the specific gene regions not routinely covered. (Unpublished Mayo method)

Genes analyzed: *ALDH18A1*, *ARG1*, *ARG2*, *ASL*, *ASS1*, *CA5A*, *CPS1*, *GLUD1*, *GLUL*, *NAGS*, *OAT*, *OTC*, *SLC25A13*, *SLC25A15*, *SLC7A7*, and *UMPS*

### PDF Report

Supplemental

### Day(s) Performed

Varies

### Report Available

21 to 35 days

### Specimen Retention Time

Whole blood: 2 weeks (if available); Extracted DNA: 3 months; Blood spots, saliva, cultured fibroblasts: 1 month

**Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

**Fees & Codes****Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

**CPT Code Information**

81443

88233-Tissue culture, skin, solid tissue biopsy (if appropriate)

88240-Cryopreservation (if appropriate)

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
UCDP	Urea Cycle Disorders Gene Panel	105267-9

Result ID	Test Result Name	Result LOINC® Value
608644	Test Description	62364-5
608645	Specimen	31208-2
608646	Source	31208-2
608647	Result Summary	50397-9
608648	Result	82939-0
608649	Interpretation	69047-9
608650	Resources	99622-3
608651	Additional Information	48767-8
608652	Method	85069-3
608653	Genes Analyzed	48018-6
608654	Disclaimer	62364-5
608655	Released By	18771-6