

Neurofibromatosis Type 1, NF1, Full Gene Analysis, Varies

### Overview

#### **Useful For**

Evaluating patients with a personal or family history suggestive of neurofibromatosis type 1 (NF1)

Establishing a diagnosis of a NF1 allowing for targeted cancer surveillance based on associated risks

Identifying genetic variants associated with increased risk for NF1 allowing for predictive testing of at-risk family members

#### **Reflex Tests**

Test Id	Reporting Name	Available Separately	Always Performed
_STR1	Comp Analysis using STR	No, (Bill only)	No
	(Bill only)		
_STR2	Add'l comp analysis w/STR	No, (Bill only)	No
	(Bill Only)		
CULFB	Fibroblast Culture for	Yes	No
	Genetic Test		
CULAF	Amniotic Fluid	Yes	No
	Culture/Genetic Test		
MATCC	Maternal Cell	Yes	No
	Contamination, B		

### **Genetics Test Information**

This test utilizes next-generation sequencing to detect single nucleotide and copy number variants in the *NF1* gene associated with neurofibromatosis type 1 (NF1). See Method Description for additional details.

Identification of a disease-causing variant may assist with diagnosis, prognosis, clinical management, familial screening, and genetic counseling for NF1

## **Testing Algorithm**

## Prenatal specimens only:

If an amniotic fluid specimen or cultured amniocytes are received, an amniotic fluid culture will be performed at an additional charge.

If a chorionic villi specimen or cultured chorionic villi are received, a fibroblast culture will be performed at an additional charge.

For any prenatal specimen that is received, maternal cell contamination testing will be performed at an additional charge.



Neurofibromatosis Type 1, NF1, Full Gene Analysis, Varies

### Skin biopsy or cultured fibroblast specimens:

For skin biopsy or cultured fibroblast specimens, a fibroblast culture will be performed at an additional charge.

## **Special Instructions**

- Molecular Genetics: Inherited Cancer Syndromes Patient Information
- Informed Consent for Genetic Testing
- Blood Spot Collection Card-Spanish Instructions
- Blood Spot Collection Card-Chinese Instructions
- Informed Consent for Genetic Testing (Spanish)
- Blood Spot Collection Instructions

#### **Method Name**

Sequence Capture and Targeted Next-Generation Sequencing (NGS) followed by Polymerase Chain Reaction (PCR) and Sanger Sequencing

#### **NY State Available**

Yes

## **Specimen**

### **Specimen Type**

Varies

## **Ordering Guidance**

For patients suspected of having neurofibromatosis 2 (NF2) or either LZTR1- or SMARCB1-related schwannomatosis, order CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies.

For a comprehensive hereditary cancer panel that includes the NF1 gene, consider ordering 1 of the following tests:

- -BRGYP / Hereditary Breast/Gynecologic Cancer Panel, Varies
- -ENDCP / Hereditary Endocrine Cancer Panel, Varies
- -HPGLP / Hereditary Paraganglioma/Pheochromocytoma Panel, Varies

Testing for the *NF1* gene as part of a customized panel is available.. For more information see CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies. To add this gene to a panel via CGPH, use the Hereditary Cancer disease state for step 1 on the <u>custom gene ordering tool</u>.

Targeted testing for familial variants (also called site-specific or known mutations testing) is available for this gene. For more information see FMTT / Familial Variant, Targeted Testing, Varies. To obtain more information about this testing option, call 800-533-1710.

## **Specimen Required**

Patient Preparation: A previous hematopoietic stem cell transplant from an allogenic donor will interfere with testing.



Neurofibromatosis Type 1, NF1, Full Gene Analysis, Varies

For information about testing patients who have received a hematopoietic stem cell transplant, call 800-533-1710.

### Submit only 1 of the following specimens:

Specimen Type: Whole blood

Container/Tube:

Preferred: Lavender top (EDTA) or yellow top (ACD)

Acceptable: Green top (Sodium heparin)

**Specimen Volume**: 3 mL Collection Instructions:

- 1. Invert several times to mix blood.
- 2. Send whole blood specimen in original tube. **Do not aliquot**.
- 3. Whole blood collected postnatal from an umbilical cord is also acceptable. See Additional Information.

Specimen Stability Information: Ambient (preferred) 4 days/Refrigerated 4 days/Frozen 4 days

#### **Additional Information:**

- 1. Specimens are preferred to be received within 4 days of collection. Extraction will be attempted for specimens received after 4 days, and DNA yield will be evaluated to determine if testing may proceed.
- 2. To ensure minimum volume and concentration of DNA are met, the requested volume must be submitted. Testing may be canceled if DNA requirements are inadequate.
- 3. For postnatal umbilical cord whole blood specimens, maternal cell contamination studies are recommended to ensure test results reflect that of the patient tested. A maternal blood specimen is required to complete maternal cell contamination studies. Order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on both the cord blood and maternal blood specimens under separate order numbers.

Specimen Type: Saliva

Patient Preparation: Patient should not eat, drink, smoke, or chew gum 30 minutes prior to collection.

Supplies:

DNA Saliva Kit High Yield (T1007) Saliva Swab Collection Kit (T786)

Container/Tube:

Preferred: High-yield DNA saliva kit

Acceptable: Saliva swab

**Specimen Volume**: 1 Tube if using T1007 or 2 swabs if using T786 **Collection Instructions**: Collect and send specimen per kit instructions.

Specimen Stability Information: Ambient (preferred) 30 days/Refrigerated 30 days

**Additional Information:** Saliva specimens are acceptable but not recommended. Due to lower quantity/quality of DNA yielded from saliva, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.

Specimen Type: Blood spot

Supplies: Card-Blood Spot Collection (Filter Paper) (T493)

Container/Tube:

Preferred: Collection card (Whatman Protein Saver 903 Paper)

Acceptable: PerkinElmer 226 filter paper or blood spot collection card

Specimen Volume: 2 to 5 Blood spots

Collection Instructions:



Neurofibromatosis Type 1, NF1, Full Gene Analysis, Varies

- 1. An alternative blood collection option for a patient older than 1 year is a fingerstick. For detailed instructions, see <a href="How to Collect a Dried Blood Spot Sample">How to Collect a Dried Blood Spot Sample</a>.
- 2. Let blood dry on the filter paper at ambient temperature in a horizontal position for a minimum of 3 hours.
- 3. Do not expose specimen to heat or direct sunlight.
- 4. Do not stack wet specimens.
- 5. Keep specimen dry

Specimen Stability Information: Ambient (preferred)/Refrigerated

#### **Additional Information:**

- 1. Blood spot specimens are acceptable but not recommended. Multiple extractions will be required to obtain sufficient yield for supplemental analysis, and there is significant risk for test failure due to insufficient DNA.
- 2. Due to lower concentration of DNA yielded from blood spot, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.
- 3. For collection instructions, see <u>Blood Spot Collection Instructions</u>
- 4. For collection instructions in Spanish, see <u>Blood Spot Collection Card-Spanish Instructions</u> (T777)
- 5. For collection instructions in Chinese, see <u>Blood Spot Collection Card-Chinese Instructions</u> (T800)

Specimen Type: Cultured fibroblasts

Source: Skin

Container/Tube: T-25 flask Specimen Volume: 2 Flasks

Collection Instructions: Submit confluent cultured fibroblast cells from a skin biopsy. Cultured cells from a prenatal

specimen will not be accepted.

Specimen Stability Information: Ambient (preferred) <24 hours/Refrigerated <24 hours

## **Additional Information:**

- 1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
- 2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks are required to culture fibroblasts before genetic testing can occur.

Specimen Type: Skin biopsy

**Supplies:** Fibroblast Biopsy Transport Media (T115)

Container/Tube: Sterile container with any standard cell culture media (eg, minimal essential media, RPMI 1640). The

solution should be supplemented with 1% penicillin and streptomycin.

**Specimen Volume:** 4-mm Punch

Specimen Stability Information: Ambient (preferred) <24 hours/Refrigerated <24 hours

### **Additional Information:**

- 1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
- 2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks are required to culture fibroblasts before genetic testing can occur.

Specimen Type: Extracted DNA

Container/Tube:



Neurofibromatosis Type 1, NF1, Full Gene Analysis, Varies

**Preferred**: Screw Cap Micro Tube, 2 mL with skirted conical base

Acceptable: Matrix tube, 1 mL

**Collection Instructions:** 

- 1. The preferred volume is at least 100 mcL at a concentration of 75 ng/mcL.
- 2. Include concentration and volume on tube.

Specimen Stability Information: Frozen (preferred) 1 year/Ambient/Refrigerated

**Additional Information**: DNA must be extracted in a CLIA-certified laboratory or equivalent and must be extracted from a specimen type listed as acceptable for this test (including applicable anticoagulants). Our laboratory has experience with Chemagic, Puregene, Autopure, MagnaPure, and EZ1 extraction platforms and cannot guarantee that all extraction methods are compatible with this test. If testing fails, one repeat will be attempted, and if unsuccessful, the test will be reported as failed and a charge will be applied. If applicable, specific gene regions that were unable to be interrogated due to DNA quality will be noted in the report.

#### **PRENATAL SPECIMENS**

**Due to its complexity, consultation with the laboratory is required** for all prenatal testing; call 800-533-1710 to speak to a genetic counselor.

Specimen Type: Amniotic fluid

Container/Tube: Amniotic fluid container

Specimen Volume: 20 mL

Specimen Stability Information: Ambient (preferred) <24 hours/Refrigerated <24 hours

**Additional Information**: Specimen will only be tested after culture.

- 1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
- 2. A separate culture charge will be assessed under CULAF / Culture for Genetic Testing, Amniotic Fluid. An additional 2 to 3 weeks are required to culture amniotic fluid before genetic testing can occur.
- 3. **All prenatal specimens must be accompanied by a maternal blood specimen;** order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on the maternal specimen.

Specimen Type: Confluent cultured amniocytes

Container/Tube: T-25 flask Specimen Volume: 2 Full flasks

**Collection Instructions:** Submit confluent cultured amniocytes from another laboratory. **Specimen Stability Information:** Ambient (preferred) <24 hours/Refrigerated <24 hours

#### **Additional Information:**

- 1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
- 2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing.
- **3.** All prenatal specimens must be accompanied by a maternal blood specimen; order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on the maternal specimen.

Specimen Type: Chorionic villi

Container/Tube: 15-mL tube containing 15 mL of transport media



Neurofibromatosis Type 1, NF1, Full Gene Analysis, Varies

Specimen Volume: 20 mg

Specimen Stability Information: Ambient (preferred) <24 hours/Refrigerated <24 hours

**Additional Information**: Specimen will only be tested after culture.

- 1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
- 2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks are required to culture fibroblasts before genetic testing can occur.
- 3. **All prenatal specimens must be accompanied by a maternal blood specimen;** order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on the maternal specimen.

Specimen Type: Cultured chorionic villi

Container/Tube: T-25 flasks Specimen Volume: 2 Full flasks

**Collection Instructions**: Submit confluent cultured cells from another laboratory

Specimen Stability Information: Ambient (preferred) <24 hours/Refrigerated <24 hours

**Additional Information:** 

- 1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
- 2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing.
- 3. **All prenatal specimens must be accompanied by a maternal blood specimen;** order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on the maternal specimen.

### **Forms**

- 1. **New York Clients-Informed consent is required.** Document on the request form or electronic order that a copy is on file. The following documents are available:
- -Informed Consent for Genetic Testing (T576)
- -Informed Consent for Genetic Testing (Spanish) (T826)
- 2. Molecular Genetics: Inherited Cancer Syndromes Patient Information Sheet (T519)
- 3. If not ordering electronically, complete, print, and send a Oncology Test Request (T729) with the specimen.

### Specimen Minimum Volume

See Specimen Required

### Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

## **Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Varies	Varies		

## Clinical & Interpretive

### **Clinical Information**



Neurofibromatosis Type 1, NF1, Full Gene Analysis, Varies

Germline variants in the *NF1* gene are associated with neurofibromatosis type 1 (NF1), an autosomal dominant hereditary tumor syndrome.(1) NF1 is characterized by many manifestations beginning in childhood, including multiple cafe au lait macules, axillary and inguinal freckling, cutaneous and plexiform neurofibromas, iris hamartomas (known as Lisch nodules), and increased lifetime risk to develop optic glioma, other brain tumors, malignant peripheral nerve sheath tumors, and breast cancer.(1) NF1 has also been associated with learning disabilities, vasculopathy, and musculoskeletal features such as osteopenia, long bone dysplasia, and scoliosis.(1) Almost half of all individuals with NF1 have no family history as their variants arose *de novo*, during gamete formation or early embryogenesis.(1)

Of note, some individuals may present with segmental/mosaic *NF1*, where an *NF1* variant may be localized to only one segment or a few segments of the body. This test may not detect mosaic *NF1*.(1)

The National Comprehensive Cancer Network provides recommendations regarding the breast cancer surveillance of adults with *NF1*.(2) Other medical management guidelines have been published by the American Academy of Pediatrics and the American College of Medical Genetics.(3,4)

## **Reference Values**

An interpretive report will be provided.

## Interpretation

All detected variants are evaluated according to American College of Medical Genetics and Genomics recommendations.(5) Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.

#### Cautions

#### Clinical Correlations:

Test results should be interpreted in the context of clinical findings, family history, and other laboratory data. Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

If testing was performed because of a clinically significant family history, it is often useful to first test an affected family member. Detection of a reportable variant in an affected family member would allow for more informative testing of at-risk individuals.

To discuss the availability of additional testing options or for assistance in the interpretation of these results, contact Mayo Clinic Laboratories genetic counselors at 800-533-1710.

## Technical Limitations:

Next-generation sequencing may not detect all types of genomic variants. In rare cases, false-negative or false-positive results may occur. The depth of coverage may be variable for some target regions; assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be considered.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. Confirmation of select reportable variants will be performed by alternate methodologies based on



Neurofibromatosis Type 1, NF1, Full Gene Analysis, Varies

internal laboratory criteria.

This test is validated to detect 95% of deletions up to 75 base pairs (bp) and insertions up to 47 bp. Deletions-insertions (delins) of 40 or more bp, including mobile element insertions, may be less reliably detected than smaller delins.

This analysis targets single and multi-exon deletions/duplications; however, in some instances single exon resolution cannot be achieved due to isolated reduction in sequence coverage or inherent genomic complexity. Balanced structural rearrangements (such as translocations and inversions) may not be detected.

This test is not designed to detect low levels of mosaicism or to differentiate between somatic mutations and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

For detailed information regarding gene-specific performance and technical limitations, see Method Description or contact a laboratory genetic counselor.

If the patient has had an allogeneic hematopoietic stem cell transplant or a recent non-leukocyte reduced blood transfusion, results may be inaccurate due to the presence of donor DNA. Call Mayo Clinic Laboratories for instructions for testing patients who have received a bone marrow transplant.

#### **Reclassification of Variants:**

Currently, it is not standard practice for the laboratory to systematically review previously classified variants on a regular basis. The laboratory encourages healthcare professionals to contact the laboratory at any time to learn how the classification of a particular variant may have changed over time. Due to broadening genetic knowledge, it is possible that the laboratory may discover new information of relevance to the patient. Should that occur, the laboratory may issue an amended report.

### Variant Evaluation:

Evaluation and categorization of variants are performed using published American College of Medical Genetics and Genomics and the Association for Molecular Pathology recommendations as a guideline. (5) Other gene-specific guidelines may also be considered. Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance. Variants classified as benign or likely benign are not reported.

Multiple in silico evaluation tools may be used to assist in the interpretation of these results. The accuracy of predictions made by in silico evaluation tools is highly dependent upon the data available for a given gene, and periodic updates to these tools may cause predictions to change over time. Results from in silico evaluation tools should be interpreted with caution and professional clinical judgment.

Rarely, incidental or secondary findings may implicate another predisposition or presence of active disease. Incidental findings may include, but are not limited to, results related to the sex chromosomes. These findings will be carefully reviewed to determine whether they will be reported.

## **Clinical Reference**



Neurofibromatosis Type 1, NF1, Full Gene Analysis, Varies

- 1. Friedman JM. Neurofibromatosis 1. In: Adam MP, Feldman J, Mirzaa GM, et al, eds. GeneReviews [Internet]. University of Washington, Seattle; 1998. Updated April 3, 2025. Accessed April 24, 2025. Available at: www.ncbi.nlm.nih.gov/books/NBK1109/
- 2. Daly MB, Pal T, Berry MP, et al. Genetic/Familial high-risk assessment: Breast, ovarian, and pancreatic, version 2.2021, NCCN Clinical Practice Guidelines in Oncology. J Natl Compr Canc Netw. 2021;19(1):77-102
- 3. Miller DT, Freedenberg D, Schorry E, et al. Health Supervision for Children With Neurofibromatosis Type 1. Pediatrics. 2019;143(5):e20190660
- 4. Stewart DR, Korf BR, Nathanson KL, Stevenson DA, Yohay K. Care of adults with neurofibromatosis type 1: a clinical practice resource of the American College of Medical Genetics and Genomics (ACMG). Genet Med. 2018;20(7):671-682
- 5. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. Genet Med. 2015;17(5):405-424

#### **Performance**

## **Method Description**

Next-generation sequencing (NGS) and/or Sanger sequencing are performed to test for the presence of variants in coding regions and intron/exon boundaries of the *NF1* gene, as well as some other regions that have known disease-causing variants. The human genome reference GRCh37/hg19 build was used for sequence read alignment. At least 99% of the bases are covered at a read depth over 30X. Sensitivity is estimated at above 99% for single nucleotide variants, above 94% for deletion-insertions (delins) less than 40 base pairs (bp), above 95% for deletions up to 75 bp and insertions up to 47 bp. NGS and/or a polymerase chain reaction-based quantitative method is performed to test for the presence of deletions and duplications in the *NF1* gene.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences.

The reference transcript for *NF1* gene is NM\_000267.3. Reference transcript numbers may be updated due to transcript re-versioning. Always refer to the final patient report for gene transcript information referenced at the time of testing.(Unpublished Mayo method)

Confirmation of select reportable variants may be performed by alternate methodologies based on internal laboratory criteria.

### PDF Report

Supplemental

### Day(s) Performed

Varies

## **Report Available**

21 days to 28 days



Neurofibromatosis Type 1, NF1, Full Gene Analysis, Varies

## **Specimen Retention Time**

Whole blood: 28 days (if available); Extracted DNA: 3 months, Saliva: 30 days (if available); Blood Spots: 1 year (if available)

## **Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

## **Fees & Codes**

## **Fees**

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

### **Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

### **CPT Code Information**

81408

88233-Tissue culture, skin, solid tissue biopsy (if appropriate)

88240-Cryopreservation (if appropriate)

88235-Amniotic fluid culture (if appropriate)

81265-Maternal cell contamination (if appropriate)

### **LOINC®** Information

Test ID	Test Order Name	Order LOINC® Value
NF1Z	NF1 Full Gene Analysis	101385-3

Result ID	Test Result Name	Result LOINC® Value
614767	Test Description	62364-5
614768	Specimen	31208-2
614769	Source	31208-2
614770	Result Summary	50397-9
614771	Result	82939-0
614772	Interpretation	69047-9
614773	Resources	99622-3
614774	Additional Information	48767-8
614775	Method	85069-3
614776	Genes Analyzed	48018-6



Neurofibromatosis Type 1, NF1, Full Gene Analysis, Varies

614777	Disclaimer	62364-5
614778	Released By	18771-6