

Ethyl Glucuronide Screen with Reflex, Random, Urine

#### Overview

#### **Useful For**

Screening for drug abuse involving alcohol

#### **Reflex Tests**

Test Id	Reporting Name	Available Separately	Always Performed
ETGC	Ethyl Glucuronide	Yes	No
	Confirmation, U		

## **Testing Algorithm**

Testing begins with a screening assay. If the screen is positive, then the liquid chromatography tandem mass spectrometry confirmation with quantification will be performed at an additional charge.

#### **Method Name**

Only orderable as part of profile. For more information see CSMEU / Controlled Substance Monitoring Enhanced Profile with Reflex, 21 Drug Classes, High Resolution Mass Spectrometry and Immunoassay Screen, Random, Urine.

**Immunoassay** 

#### **NY State Available**

Yes

## Specimen

# Specimen Type

Urine

#### Specimen Required

Only orderable as part of profile. For more information see CSMEU / Controlled Substance Monitoring Enhanced Profile with Reflex, 21 Drug Classes, High Resolution Mass Spectrometry and Immunoassay Screen, Random, Urine.

Container/Tube: Plastic, 60-mL urine bottle

**Specimen Volume:** 5 mL **Collection Instructions:** 

1. Collect a random urine specimen.

2. No preservative.

## **Specimen Minimum Volume**



Ethyl Glucuronide Screen with Reflex, Random, Urine

2.5 mL

# **Reject Due To**

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

# **Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	28 days	
	Ambient	28 days	
	Frozen	28 days	

# Clinical & Interpretive

#### Clinical Information

This test uses immunoassay reagents designed to produce a negative result when no drugs are present in a natural (ie, unadulterated) urine specimen; the assay is designed to have a high true-negative rate. Like all immunoassays, it can have a false-positive rate due to cross-reactivity with natural chemicals and drugs other than those they were designed to detect. The immunoassay also has a false-negative rate to the antibody's ability to cross-react with different drugs in the class being screened.

Ethyl glucuronide is a direct metabolite of ethanol formed by enzymatic conjugation of ethanol with glucuronic acid. Alcohol in urine is normally detected for only a few hours, whereas ethyl glucuronide can be detected in the urine for 1 to 3 days.

#### **Reference Values**

Only orderable as part of profile. For more information see CSMEU / Controlled Substance Monitoring Enhanced Profile with Reflex, 21 Drug Classes, High Resolution Mass Spectrometry and Immunoassay Screen, Random, Urine.

Negative

Screening cutoff concentration: Ethyl glucuronide: 500 ng/mL

#### Interpretation

This assay only provides a preliminary analytical test result. A more specific alternative method (ie, liquid chromatography tandem mass spectrometry) must be used to obtain a confirmed analytical result. A positive result using the ethyl glucuronide screen indicates only the potential presence of ethyl glucuronide and does not necessarily correlate with the extent of physiological and psychological effects.

# **Cautions**

Care should be taken when interpreting results since there are many factors (eg, fluid intake and other biologic factors) that may influence a urine test result. It is possible that substances other than those investigated in the specificity study



Ethyl Glucuronide Screen with Reflex, Random,
Urine

may interfere with the test and cause false-positive or negative results.

#### **Clinical Reference**

- 1. Schmitt G, Aderjan R, Keller T, Wu M. Ethyl glucuronide: an unusual ethanol metabolite in humans. Synthesis, analytical data, and determination in serum and urine. J Anal Toxicol. 1995;19(2):91-94
- 2. Dahl H, Stephanson N, Beck O, Helander A. Comparison of urinary excretion characteristics of ethanol and ethyl glucuronide. J Anal Toxicol. 2002;26(4):201-204. doi:10.1093/jat/26.4.201
- 3. Wurst FM, Skipper GE, Weinmann W. Ethyl glucuronide--the direct ethanol metabolite on the threshold from science to routine use. Addiction. 2003;98 (Suppl 2):51-61. doi:10.1046/j.1359-6357.2003.00588.x
- 4. Zimmer H, Schmitt G, Aderjan R. Preliminary immunochemical test for the determination of ethyl glucuronide in serum and urine: comparison of screening method results with gas chromatography-mass spectrometry. J Anal Toxicol. 2002;26(1):11-16. doi:10.1093/jat/26.1.11
- 5. Weinmann W, Schaefer P, Thierauf A, Schreiber A, Wurst FM. Confirmatory analysis of ethyl glucuronide in urine by liquid chromatography/electrospray ionization/tandem mass spectrometry according to forensic guidelines. J Am Soc Mass Spectrom. 2004;15(2):188-193. doi:10.1016/j.jasms.2003.10.010
- 6. Langman LJ, Bechtel LK, Holstege CP. Clinical toxicology. In: Rifai N, Chiu RWK, Young I, Burnham CAD, Wittwer CT, eds. Tietz Textbook of Laboratory Medicine. 7th ed. Elsevier; 2023:chap 43

#### **Performance**

### **Method Description**

This assay is a homogeneous enzyme-linked immunosorbent assay technique. The assay will be performed semiquantitatively. The assay is based on competition between free drug in the urine sample, and a drug labeled with the enzyme glucose-6-phosphate dehydrogenase for a fixed amount of specific antibody binding sites. Active enzyme converts nicotinamide adenine dinucleotide (NAD[+]) to NADH, which results in an absorbance change that can be measured spectrophotometrically at 340 nm.(Package insert: ETG. Immunalysis; 04/2019)

## **PDF Report**

No

### Day(s) Performed

Monday through Saturday

# **Report Available**

Same/1 to 2 days

#### Specimen Retention Time

14 days

# **Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Superior Drive



Ethyl Glucuronide Screen with Reflex, Random, Urine

# **Fees & Codes**

## **Fees**

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

#### **Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

## **CPT Code Information**

80307

#### **LOINC®** Information

Test ID	Test Order Name	Order LOINC® Value
ETGSR	Ethyl Glucuronide Scrn w/Reflex, U	58375-7

Result ID	Test Result Name	Result LOINC® Value
616033	Ethyl Glucuronide Screen, U	58375-7