

Overview

Useful For

Reporting an end titer result for tripartite motif-containing protein 46 (TRIM46)-IgG in serum specimens

Evaluation of an autoimmune/paraneoplastic neurological syndrome among patients presenting with cerebellar ataxia, encephalitis, or encephalomyelitis.

Testing Algorithm

If the indirect immunofluorescence (IFA) pattern suggests tripartite motif-containing protein 46 (TRIM46) IgG, then the TRIM46 antibody cell-binding assay (CBA) and TRIM46 antibody IFA titer will be performed at an additional charge.

Method Name

Only orderable as a reflex. For more information see:

ENS2 / Encephalopathy, Autoimmune/Paraneoplastic Evaluation, Serum

DMS2 / Dementia, Autoimmune/Paraneoplastic Evaluation, Serum

EPS2 / Epilepsy, Autoimmune/Paraneoplastic Evaluation, Serum

MDS2 / Movement Disorder, Autoimmune/Paraneoplastic Evaluation, Serum

MAS1 / Myelopathy, Autoimmune/Paraneoplastic Evaluation, Serum

Indirect Immunofluorescence Assay (IFA)

NY State Available

Yes

Specimen

Specimen Type

Serum

Specimen Required

Only orderable as a reflex. For more information see:

ENS2 / Encephalopathy, Autoimmune/Paraneoplastic Evaluation, Serum

DMS2 / Dementia, Autoimmune/Paraneoplastic Evaluation, Serum

EPS2 / Epilepsy, Autoimmune/Paraneoplastic Evaluation, Serum

MDS2 / Movement Disorder, Autoimmune/Paraneoplastic Evaluation, Serum

MAS1 / Myelopathy, Autoimmune/Paraneoplastic Evaluation, Serum

Reject Due To

| | |
|-----------------|--------|
| Gross hemolysis | Reject |
| Gross lipemia | Reject |
| Gross icterus | Reject |

Specimen Stability Information

| Specimen Type | Temperature | Time | Special Container |
|---------------|--------------------------|----------|-------------------|
| Serum | Refrigerated (preferred) | 28 days | |
| | Ambient | 72 hours | |
| | Frozen | 28 days | |

Clinical & Interpretive

Clinical Information

Tripartite motif-containing protein 46 (TRIM46-IgG) is a marker of an autoimmune neurological disorder commonly associated with underlying malignancy. Patients commonly present with cerebellar ataxia and neoplasms frequently of neuroendocrine lineage.

Reference Values

Only orderable as a reflex. For more information see:

ENS2 / Encephalopathy, Autoimmune/Paraneoplastic Evaluation, Serum

DMS2 / Dementia, Autoimmune/Paraneoplastic Evaluation, Serum

EPS2 / Epilepsy, Autoimmune/Paraneoplastic Evaluation, Serum

MDS2 / Movement Disorder, Autoimmune/Paraneoplastic Evaluation, Serum

MAS1 / Myelopathy, Autoimmune/Paraneoplastic Evaluation, Serum

<1:240

Interpretation

A positive result is consistent with a tripartite motif-containing protein 46 (TRIM46-IgG) associated autoimmune disease of the central nervous system. A paraneoplastic cause should be considered.

Cautions

A negative result does not exclude the presence of neurological autoimmunity or cancer. The use of immunotherapy prior to sample collection may negatively impact the sensitivity of this assay.

Clinical Reference

1. van Coevorden-Hameete MH, van Beuningen SFB, Perrenoud M, et al. Antibodies to TRIM46 are associated with paraneoplastic neurological syndromes. *Ann Clin Tran Neurol.* 2017;4(9):680-686. doi:10.1002/acn3.396
2. Shams'ili S, de Leeuw B, Hulsenboom E, Jaarsma D, Smitt PS. A new paraneoplastic encephalomyelitis autoantibody

reactive with the axon initial segment. *Neurosci Lett.* 2009;467(2):169-172. doi:10.1016/j.neulet.2009.10.031

3. Valencia-Sanchez C, Knight AM, Hammami B, et al. TRIM46 autoantibody: expanded neurological phenotype and oncological associations (1657). *Neurology.* 2021;96(15 Supplement). doi:10.1212/WNL.96.15_supplement.1657

Performance

Method Description

The patient's specimen is tested by a standardized immunofluorescence assay that uses a composite frozen section of mouse cerebellum, kidney, and gut tissues. After incubation with the specimen and washing, fluorescein-conjugated goat-antihuman IgG is applied. Neuron-specific autoantibodies are identified by their characteristic fluorescence staining patterns. Specimens that are scored positive for any neuronal nuclear or cytoplasmic autoantibody are titrated. Interference by coexisting non-neuron-specific autoantibodies can usually be eliminated by serologic absorption.(Honorat JA, Komorowski L, Josephs KA, et al. IgLON5 antibody: Neurological accompaniments and outcomes in 20 patients. *Neurol Neuroimmunol Neuroinflamm.* 2017;4[5]:e385. Published 2017 Jul 18. doi:10.1212/NXI.0000000000000385)

PDF Report

No

Day(s) Performed

Monday through Sunday

Report Available

5 to 10 days

Specimen Retention Time

2 days

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA

requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

86256

LOINC® Information

| Test ID | Test Order Name | Order LOINC® Value |
|---------|------------------------|--------------------|
| T46TS | TRIM46 Ab IFA Titer, S | 105527-6 |

| Result ID | Test Result Name | Result LOINC® Value |
|-----------|------------------------|---------------------|
| 616447 | TRIM46 Ab IFA Titer, S | 105527-6 |