

Hypobetalipoproteinemia Gene Panel, Varies

### Overview

#### Useful For

Providing a genetic evaluation for patients with a personal or family history suggestive of familial hypobetalipoproteinemia

#### **Reflex Tests**

Test Id	Reporting Name	Available Separately	Always Performed
CULFB	Fibroblast Culture for	Yes	No
	Genetic Test		

#### **Genetics Test Information**

This test utilizes next-generation sequencing to detect single nucleotide and copy number variants in 5 genes associated with familial hypobetalipoproteinemia: *ANGPTL3, APOB, MTTP, PCSK9,* and *SAR1B*. See <u>Targeted Genes and</u> <u>Methodology Details for Hypobetalipoproteinemia Gene Panel</u> and Method Description for additional details.

Identification of a disease-causing variant may assist with diagnosis, prognosis, clinical management, familial screening, and genetic counseling for familial hypobetalipoproteinemia.

Prior Authorization is available for this assay.

#### **Testing Algorithm**

**Skin biopsy:** For skin biopsy or cultured fibroblast specimens, a fibroblast culture will be performed at an additional charge. If viable cells are not obtained, the client will be notified.

#### **Special Instructions**

- Informed Consent for Genetic Testing
- Informed Consent for Genetic Testing (Spanish)
- Hereditary Dyslipidemia Patient Information
- Targeted Genes and Methodology Details for Hypobetalipoproteinemia Gene Panel
- Hypobetalipoproteinemia Gene Panel (HYPBG) Prior Authorization Ordering Instructions

#### Method Name

Sequence Capture and Targeted Next-Generation Sequencing followed by Polymerase Chain Reaction (PCR) and Sanger Sequencing

#### NY State Available

Yes

#### Specimen



Hypobetalipoproteinemia Gene Panel, Varies

## **Specimen Type**

Varies

## **Ordering Guidance**

This gene panel contains genes in common with HCHLG / Hypercholesterolemia Gene Panel, Varies. These tests should not be ordered concurrently as they assist in diagnosing conflicting disorders. For low levels of cholesterol (hypocholesterolemia), order this test. For high levels of cholesterol (hypercholesterolemia), order HCHLG.

Customization of this panel and single gene analysis for any gene present on this panel are available. For more information see CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies. To modify this panel via CGPH, use the Cardiovascular/Connective Tissue/Dyslipidemia/Cerebrovascular/Primary Ciliary Dyskinesia disease state for step 1 on the <u>Custom Gene Ordering Tool</u>.

Targeted testing for familial variants (also called site-specific or known mutations testing) is available for the genes on this panel. See FMTT / Familial Variant, Targeted Testing, Varies. To obtain more information about this testing option, call 800-533-1710.

## **Necessary Information**

<u>Prior Authorization</u> is available, **but not required**, for this test. If proceeding with the prior authorization process, submit the required form with the specimen.

## Specimen Required

**Patient Preparation:** A previous hematopoietic stem cell transplant from an allogenic donor will interfere with testing. For instructions for testing patients who have received a hematopoietic stem cell transplant, call 800-533-1710.

### Submit only 1 of the following specimens:

Specimen Type: Whole blood
Container/Tube:
Preferred: Lavender top (EDTA) or yellow top (ACD)
Acceptable: Green top (Sodium heparin)
Specimen Volume: 3 mL
Collection Instructions:

Invert several times to mix blood.
Send whole blood specimen in original tube. Do not aliquot.

Specimen Stability Information: Ambient (preferred) 4 days/Refrigerated 4 days/Frozen 4 days

## Additional Information:

1. Specimens are preferred to be received within 4 days of collection. Extraction will be attempted for specimens received after 4 days, and DNA yield will be evaluated to determine if testing may proceed.

2. To ensure minimum volume and concentration of DNA is met, the requested volume must be submitted. Testing may be canceled if DNA requirements are inadequate.

Specimen Type: Cord blood Container/Tube: Preferred: Lavender top (EDTA) or yellow top (ACD)





Hypobetalipoproteinemia Gene Panel, Varies

Acceptable: Green top (Sodium heparin)

Specimen Volume: 3 mL

**Collection Instructions:** 

1. Invert several times to mix blood.

2. Send cord blood specimen in original tube. Do not aliquot.

Specimen Stability Information: Ambient (preferred) 4 days/Refrigerated 4 days/Frozen 4 days

### Additional Information:

1. Specimens are preferred to be received within 4 days of collection. Extraction will be attempted for specimens received after 4 days, and DNA yield will be evaluated to determine if testing may proceed.

2. To ensure minimum volume and concentration of DNA is met, the requested volume must be submitted. Testing may be canceled if DNA requirements are inadequate.

3. While a properly collected cord blood sample may not be at risk for maternal cell contamination, unanticipated complications may occur during collection. Therefore, maternal cell contamination studies are recommended to ensure the test results reflect that of the patient tested and are available at an additional charge. Order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on the maternal specimen.

Specimen Type: Saliva

Patient Preparation: Patient should not eat, drink, smoke, or chew gum 30 minutes prior to collection.

Supplies: Saliva Collection Kit (T786)

Specimen Volume: 2 Swabs

**Collection Instructions**: Collect and send specimen per kit instructions.

Specimen Stability Information: Ambient (preferred) 30 days/Refrigerated 30 days

Additional information: Saliva specimens are acceptable but not recommended. Due to lower quantity/quality of DNA yielded from saliva, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.

### Specimen Type: Cultured fibroblasts

Source: Skin

Container/Tube: T-25 flask

Specimen Volume: 2 Flasks

**Collection Instructions**: Submit confluent cultured fibroblast cells from a skin biopsy. Cultured cells from a prenatal specimen will not be accepted.

Specimen Stability Information: Ambient (preferred) <24 hours/Refrigerated <24 hours

### Additional Information:

1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.

2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks are required to culture fibroblasts before genetic testing can occur.

Specimen Type: Tissue biopsy
Supplies: Hank's Solution (T132)
Container/Tube: Sterile container with sterile Hank's balanced salt solution, Ringer's solution, or normal saline.
Specimen Volume: 0.5 to 3 cm(3) or larger
Specimen Stability Information: Ambient (preferred) <24 hours/Refrigerated <24 hours</li>



### Hypobetalipoproteinemia Gene Panel, Varies

#### Additional Information:

1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.

2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks are required to culture fibroblasts before genetic testing can occur.

#### Specimen Type: Blood spot

Supplies: Card-Blood Spot Collection (Filter Paper) (T493)

#### Container/Tube:

Preferred: Collection card (Whatman Protein Saver 903 Paper)

Acceptable: PerkinElmer 226 filter paper or blood spot collection card

Specimen Volume: 2 to 5 Blood spots

#### Collection Instructions:

1. An alternative blood collection option for a patient older than 1 year is a fingerstick. For detailed instructions, see <u>How to Collect a Dried Blood Spot Sample</u>.

- 2. Let blood dry on the filter paper at ambient temperature in a horizontal position for a minimum of 3 hours.
- 3. Do not expose specimen to heat or direct sunlight.
- 4. Do not stack wet specimens.
- 5. Keep specimen dry.

### Specimen Stability Information:: Ambient (preferred)/Refrigerated

#### Additional Information:

1. Blood spot specimens are acceptable but not recommended. Due to lower quantity/quality of DNA yielded from blood spots, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.

2. Due to lower concentration of DNA yielded from blood spot, it is possible that additional specimen may be required to complete testing.

3. For collection instructions, see <u>Blood Spot Collection Instructions</u>

- 4. For collection instructions in Spanish, see <u>Blood Spot Collection Card-Spanish Instructions</u> (T777)
- 5. For collection instructions in Chinese, see <u>Blood Spot Collection Card-Chinese Instructions</u> (T800)

### Specimen Type: Extracted DNA

### **Container/Tube**:

Preferred: Screw Cap Micro Tube, 2 mL with skirted conical base

Acceptable: Matrix tube, 1 mL

### Collection Instructions:

- 1. The preferred volume is at least 100 mcL at a concentration of 75 ng/mcL.
- 2. Include concentration and volume on tube.
- Specimen Stability Information: Frozen (preferred) 1 year/Ambient/Refrigerated

Additional Information: DNA must be extracted in a CLIA-certified laboratory or equivalent and must be extracted from a specimen type listed as acceptable for this test (including applicable anticoagulants). Our laboratory has experience with Chemagic, Puregene, Autopure, MagnaPure, and EZ1 extraction platforms and cannot guarantee that all extraction methods are compatible with this test. If testing fails, one repeat will be attempted, and if unsuccessful, the test will be reported as failed and a charge will be applied. If applicable, specific gene regions that were unable to be interrogated due to DNA quality will be noted in the report.



Hypobetalipoproteinemia Gene Panel, Varies

#### Forms

1. New York Clients-Informed consent is required. Document on the request form or electronic order that a copy is on file.

The following documents are available:

-Informed Consent for Genetic Testing (T576)

-Informed Consent for Genetic Testing (Spanish) (T826)

2. <u>Hereditary Dyslipidemia Patient Information</u>

3. <u>Hypobetalipoproteinemia Gene Panel (HYPBG) Prior Authorization Ordering Instructions</u>

4. If not ordering electronically, complete, print, and send a <u>Cardiovascular Test Request</u> (T724) with the specimen.

### **Specimen Minimum Volume**

See Specimen Required

### Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

#### **Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Varies	Varies		

#### Clinical & Interpretive

#### **Clinical Information**

Monogenic causes of hypobetalipoproteinemia include familial hypobetalipoproteinemia (FHBL), abetalipoproteinemia (ABL), chylomicron retention disease (CMRD), loss-of-function variants in *PCSK9*, and familial combined hypolipidemia (FCH).

Familial hypobetalipoproteinemia is the most common form of hypobetalipoproteinemia and is characterized by apolipoprotein B (ApoB) levels below the 5th percentile and low-density lipoprotein cholesterol (LDL-C) concentrations in the range of 20 mg/dL to 50 mg/dL. FHBL displays codominant inheritance, whereby heterozygous individuals may be asymptomatic or have mild disease, and (rare) compound heterozygous and homozygous individuals develop more severe, early-onset disease. In cases of mild to moderate FHBL with little or no liver involvement, prognosis is good and, in fact, may be associated with increased longevity. In severe disease, symptoms include fatty liver, which may progress to cirrhosis over time, symptoms of fat malabsorption, failure to thrive, and neurological and ocular dysfunction. FHBL is most commonly due to disease-causing truncating variants in the *APOB* gene resulting in reduced or nonfunctional protein.

Abetalipoproteinemia is a rare (<1:1,000,000) condition characterized by triglyceride concentrations of less than 30 mg/dL, cholesterol concentrations of less than 30 mg/dL, and undetectable LDL and ApoB levels. Clinical presentation is similar to that described above for compound heterozygous and homozygous FHBL. ABL displays autosomal recessive inheritance and is caused by compound heterozygous or homozygous disease-causing variants in the *MTTP* gene.

Chylomicron retention disease is a rare lipid malabsorption syndrome that typically presents in young infants with diarrhea, steatorrhea, abdominal distention, and failure to thrive. Laboratory findings include LDL-C and high-density



Hypobetalipoproteinemia Gene Panel, Varies

lipoprotein cholesterol-C reduction of approximately 50% with normal triglyceride concentrations. CMRD displays autosomal recessive inheritance and is caused by compound heterozygous or homozygous disease-causing variants in the *SAR1B* gene.

Heterozygous loss-of-function variants in the *PCSK9* gene are associated with mild to moderate reduction in LDL-C and normal health with significantly lower prevalence of atherosclerotic heart disease. Rare individuals with biallelic loss-of-function variants in *PCSK9* have been reported with extremely low levels of LDL-C (approximately 15 mg/dL), normal health and reproductive capacity, and no evidence of neurologic or cognitive dysfunction. Notably, heterozygous gain-of-function variants in *PCSK9* are associated with familial hypercholesterolemia.

Finally, FCH is a very rare condition of panhypolipidemia associated with normal health and significantly lower prevalence of atherosclerotic heart disease. This condition is caused by loss-of-function variants in the *ANGPTL3* gene. Similar to FHBL and *PCSK9*, FCH displays codominant inheritance, with heterozygotes having normal HDL-C and LDL-C concentrations that are below the 25th percentile, while compound heterozygous and homozygous individuals display significant reductions in HDL-C levels as well.

### **Reference Values**

An interpretive report will be provided.

### Interpretation

All detected variants are evaluated according to American College of Medical Genetics and Genomics recommendations.(1) Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.

## Cautions

### Clinical Correlations:

Test results should be interpreted in the context of clinical findings, family history, and other laboratory data. Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

If testing was performed because of a clinically significant family history, it is often useful to first test an affected family member. Detection of a reportable variant in an affected family member would allow for more informative testing of at-risk individuals.

To discuss the availability of additional testing options or for assistance in the interpretation of these results, contact the Mayo Clinic Laboratories genetic counselors at 800-533-1710.

### Technical Limitations:

Next-generation sequencing may not detect all types of genomic variants. In rare cases, false-negative or false-positive results may occur. The depth of coverage may be variable for some target regions; assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be considered.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. Confirmation of select reportable variants will be performed by alternate methodologies based on



### Hypobetalipoproteinemia Gene Panel, Varies

#### internal laboratory criteria.

This test is validated to detect 95% of deletions up to 75 base pairs (bp) and insertions up to 47 bp. Deletions-insertions (delins) of 40 or more bp, including mobile element insertions, may be less reliably detected than smaller delins.

#### Deletion/Duplication Analysis:

This analysis targets single and multi-exon deletions/duplications; however, in some instances, single exon resolution cannot be achieved due to isolated reduction in sequence coverage or inherent genomic complexity. Balanced structural rearrangements (such as translocations and inversions) may not be detected.

This test is not designed to detect low levels of mosaicism or to differentiate between somatic mutations and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

Genes may be added or removed based on updated clinical relevance. For the most up to date list of genes included in this test and detailed information regarding gene specific performance and technical limitations, see Method Description or contact a laboratory genetic counselor.

If the patient has had an allogeneic hematopoietic stem cell transplant or a recent blood transfusion, results may be inaccurate due to the presence of donor DNA. Call Mayo Clinic Laboratories for instructions for testing patients who have received a bone marrow transplant.

#### **Reclassification of Variants:**

Currently, it is not standard practice for the laboratory to systematically review previously classified variants on a regular basis. The laboratory encourages healthcare professionals to contact the laboratory at any time to learn how the classification of a particular variant may have changed over time.

#### Variant Evaluation:

Evaluation and categorization of variants are performed using published American College of Medical Genetics and Genomics and the Association for Molecular Pathology recommendations as a guideline.(1) Other gene-specific guidelines may also be considered. Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance. Variants classified as benign or likely benign are not reported.

Multiple in silico evaluation tools may be used to assist in the interpretation of these results. The accuracy of predictions made by in silico evaluation tools is highly dependent upon the data available for a given gene, and periodic updates to these tools may cause predictions to change over time. Results from in silico evaluation tools should be interpreted with caution and professional clinical judgment.

Rarely, incidental or secondary findings may implicate another predisposition or presence of active disease. Incidental findings may include, but are not limited to, results related to the sex chromosomes. These findings will be carefully reviewed to determine whether they will be reported.

### **Clinical Reference**

1. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus



Hypobetalipoproteinemia Gene Panel, Varies

recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. Genet Med. 2015;17(5):405-424

2. Shapiro MD and Feingold KR. Monogenic disorders causing hypobetalipoproteinemia. In: Feingold KR, Anawalt B, Boyce A, et al, eds. Endotext [Internet]. MDTest.com, Inc; 2000. Updated January 22, 2024. Accessed March 31, 2025. Available at www.ncbi.nlm.nih.gov/books/NBK326744/

## Performance

## Method Description

Next-generation sequencing (NGS) and/or Sanger sequencing is performed to test for the presence of variants in coding regions and intron/exon boundaries of the genes analyzed, as well as some other regions that have known disease-causing variants. The human genome reference GRCh37/hg19 build was used for sequence read alignment. At least 99% of the bases are covered at a read depth over 30X. Sensitivity is estimated at above 99% for single nucleotide variants, above 94% for deletions-insertions (delins) less than 40 base pairs (bp), above 95% for deletions up to 75 bp and insertions up to 47 bp. NGS and/or a polymerase chain reaction-based quantitative method is performed to test for the presence of deletions and duplications in the genes analyzed. Confirmation of select reportable variants may be performed by alternate methodologies based on internal laboratory criteria.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. See <u>Targeted Genes and Methodology Details for Hypobetalipoproteinemia Gene Panel</u> for details regarding the targeted genes analyzed for each test and specific gene regions not routinely covered.(Unpublished Mayo method)

Genes analyzed: ANGPTL3, APOB, MTTP, PCSK9, and SAR1B

PDF Report Supplemental

Day(s) Performed Varies

Report Available 21 to 28 days

**Specimen Retention Time** Whole blood: 28 days (if available); Salvia: 30 days (if available); Extracted DNA: 3 months; Blood spots: 1 year Whole

**Performing Laboratory Location** Mayo Clinic Laboratories - Rochester Main Campus

## Fees & Codes



Hypobetalipoproteinemia Gene Panel, Varies

#### Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact Customer Service.

### **Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

### **CPT Code Information**

81406
81407
81479
81479 (if appropriate for government payers)
88233-Tissue culture, skin, solid tissue biopsy (if appropriate)
88240-Cryopreservation (if appropriate)

### **Prior Auhtorization**

Insurance preauthorization is available for this testing; forms are available.

Patient financial assistance may be available to those who qualify. Patients who receive a bill from Mayo Clinic Laboratories will receive information on eligibility and how to apply.

### LOINC<sup>®</sup> Information

Test ID	Test Order Name	Order LOINC <sup>®</sup> Value
НҮРВБ	Hypobetalipoproteinemia Gene	51966-0
	Panel	
Result ID	Test Result Name	Result LOINC <sup>®</sup> Value
617310	Test Description	62364-5
617311	Specimen	31208-2
617312	Source	31208-2
617313	Result Summary	50397-9
617314	Result	82939-0
617315	Interpretation	69047-9
617316	Additional Results	82939-0
617317	Resources	99622-3
617318	Additional Information	48767-8
617319	Method	85069-3
617320	Genes Analyzed	48018-6
617321	Disclaimer	62364-5
617322	Released By	18771-6