

## Overview

### Useful For

Evaluation of patients with suspected Brazil nut allergy to component Ber e 1

### Testing Algorithm

If the Brazil nut specific total IgE result is 0.10 kU/L or more, then the Brazil nut component (Ber e 1) test is performed at an additional charge.

### Method Name

Only orderable as a reflex. For more information see BRAZR / Brazil Nut, IgE, with Reflex to Brazil Nut Component, IgE, Serum.

Fluorescent Enzyme Immunoassay (FEIA)

### NY State Available

Yes

## Specimen

### Specimen Type

Serum

### Specimen Required

Only orderable as a reflex. For more information see BRAZR / Brazil Nut, IgE, with Reflex to Brazil Nut Component, IgE, Serum.

**Supplies:** Sarstedt Aliquot Tube, 5 mL (T914)

**Collection Container/Tube:**

**Preferred:** Serum gel

**Acceptable:** Red top

**Submission Container/Tube:** Plastic vial

**Specimen Volume:** 0.6 mL Serum

**Collection Instructions:** Centrifuge and aliquot serum into a plastic vial.

### Specimen Minimum Volume

Serum: 0.4 mL

### Reject Due To

Gross hemolysis	OK
-----------------	----

Gross lipemia	OK
Gross icterus	OK

## Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	14 days	
	Frozen	90 days	

## Clinical & Interpretive

### Clinical Information

Allergies to tree nuts are relatively prevalent and can result in severe reactions. The main culprits in tree nut allergies include walnut, almond, pistachio, cashew, pecan, hazelnut, macadamia, Brazil nut, and pine nuts. Tree nut allergy often appears in young children and estimates of prevalence range from 0.1% to greater than 5% of the population, dependent on geographical region.

In the case of nut-induced allergic reactions, as with many other foods, symptoms usually present within minutes of ingestion. Over 80% of reactions to tree nuts involve allergy related respiratory symptoms. Tree nut allergies are one of the most dangerous types of allergic reaction with 20% to 40% of cases of related anaphylaxis, and 70% to 90% of fatalities attributable to nut exposure (including peanut exposure).

Allergy to Brazil nut has reported within the United States population. Among those suffering from tree-nut-allergic individuals, the prevalence of Brazil nut allergy is estimated to be 10% to 20%. Brazil nut allergy occurs primarily through oral ingestion. Following oral exposure, allergy may be associated with systemic reactions, including respiratory and urticaria, occasionally resulting in anaphylaxis.

Ber e 1 is an abundant, heat and digestion resistant, storage protein component that is associated with systemic reactions to Brazil nuts. This major allergen component has been found to correlate with allergic symptoms. Exposure of the Ber e 1 at 100 degrees C for 20 minutes did not reduce the potential allergenicity of the molecule.

Immunological cross-reactivity has been reported between Brazil nut, hazelnut, cashew, pistachio, and almond, although cross reactivity with walnut, peanut, and coconut has also been reported. There is considerable homology between the 2S albumin of Brazil nut (Ber e 1) and other plant species, such as cottonseed, sunflower, rapeseed, castor bean, and sesame. Positive antibodies to total Brazil nuts may occur in cases of allergy to other Brazil nut storage proteins, profilins, or in the presence of cross-reacting carbohydrate determinants.

### Reference Values

Only orderable as a reflex. For more information see BRAZR / Brazil Nut, IgE, with Reflex to Brazil Nut Component, IgE, Serum

Class	IgE kU/L	Interpretation
0	<0.10	Negative
0/1	0.10-0.34	Borderline/Equivocal

1	0.35-0.69	Equivocal
2	0.70-3.49	Positive
3	3.50-17.4	Positive
4	17.5-49.9	Strongly positive
5	50.0-99.9	Strongly positive
6	> or =100	Strongly positive

Concentrations of 0.70 kU/L or more (class 2 and above) will flag as abnormally high.

Reference values apply to all ages.

### Interpretation

When detectable total Brazil nut IgE antibody is present (> or =0.10 IgE kUa/L), additional specific component IgE antibody testing will be performed. If a potential specific allergenic Brazil nut component IgE is detectable (> or =0.10 IgE kUa/L), an interpretive report will be provided.

When the sample is negative for total Brazil nut IgE antibody (<0.10 IgE kUa/L), further testing for specific Brazil nut component IgE antibodies will not be performed. A negative IgE result for total Brazil nut antibody may indicate a lack of sensitization to the potential Brazil nut allergenic component.

### Cautions

Clinical correlation of results from in vitro IgE testing with patient history of allergic or anaphylactic responses to Brazil nuts is recommended.

Negative results for IgE antibodies against Brazil nut extract or allergenic components do not completely exclude the possibility of clinically relevant allergic responses upon exposure.

Positive results for IgE to Brazil nuts or any potential Brazil nut allergenic components are not diagnostic for allergy and only indicate patient may be sensitized to Brazil nuts or a cross-reactive allergen.

Testing for IgE antibodies may not be useful in patients previously treated with immunotherapy to determine if residual clinical sensitivity exists or in patients whose medical management does not depend upon the identification of allergen specificity.

False-positive results for IgE antibodies may occur in patients with markedly elevated serum IgE (>2500 kU/L) due to nonspecific binding to allergen solid phases.

Cross-reacting carbohydrate determinants may also result in positive total Brazil nut specific IgE testing.

### Clinical Reference

1. Salo PM, Arbes SJ Jr, Jaramillo R, et al. Prevalence of allergic sensitization in the United States: results from the National Health and Nutrition Examination Survey (NHANES) 2005-2006. *J Allergy Clin Immunol.* 2014;134(2):350-359. doi:10.1016/j.jaci.2013.12.1071
2. Wasserman S, Watson W. Food allergy. *Allergy Asthma Clin Immuno.* 2011;7 Suppl 1(Suppl 1):S7
3. Abrams EM, Sicherer SH. Diagnosis and management of food allergy. *CMAJ.* 2016;188(15):1087-1093. doi:10.1503/cmaj.160124

4. Weinberger T, Sicherer S. Current perspectives on tree nut allergy: a review. *J Asthma Allergy*. 2018;11:41-51. doi:10.2147/JAA.S141636
5. Lomas JM, Jarvinen KM. Managing nut-induced anaphylaxis: challenges and solutions. *J Asthma Allergy*. 2015; 8:115-123. doi:10.2147/JAA.S89121
6. Maloney JM, Rudengren M, Ahlstedt S, Bock SA, Sampson HA. The use of serum-specific IgE measurements for the diagnosis of peanut, tree nut, and seed allergy. *J Allergy Clin Immunol*. 2008;122(1):145-151. doi:10.1016/j.jaci.2008.04.014
7. Sicherer SH, Burks AW, Sampson HA. Clinical features of acute allergic reactions to peanut and tree nuts in children. *Pediatrics*. 1998;102(1):e6. doi:10.1542/peds.102.1.e6
8. Crespo JF, James JM, Fernandez C, Rodriguez J. Food allergy: Nuts and tree nuts. *Br J Nutr*. 2006;96 Suppl 2:S95-102
9. Yang L, Clements S, Joks R. A retrospective study of peanut and tree nut allergy: Sensitization and correlations with clinical manifestations. [published online ahead of print, 2015 Feb 27]. *Allergy Rhinol (Providence)*. 2015;doi:10.2500/ar.20105.6.0108
10. Masthoff L, Hoff R, Verhoeckx KC, et al. A systematic review of the effect of thermal processing on the allergenicity of tree nuts. *Allergy*. 2013;3;68(8):983-993. doi:10.1111/all.12185
11. Borja JM, Bartolome B, Gomez E, Galindo PA, Feo F. Anaphylaxis from Brazil nut. *Allergy*. 1999;54(9):1007-1008. doi:10.1034/j.1398-9995.1999.00152.x
12. Mazokopakis EE, Lontiris MI. Commentary: Health concerns of Brazil nut consumption. *J Altern Complement Med*. 2018;24(1):3-6. doi:10.1089/acm.2017.0159
13. McWilliam V, Koplin J, Lodge C, Tang M, Dharmage S, Allen K. The prevalence of tree nut allergy: A systematic review. *Curr Allergy Asthma Rep*. 2015;15(9):54. doi:10.1007/s11882-015-0555-8
14. Rayes H, Raza AA, Williams A, Matthews S, Arshad SH. Specific IgE to recombinant protein (Ber e 1) for the diagnosis of Brazil nut allergy. *Clin Exp Allergy*. 2016;46(4):654-656. doi:10.1111/cea.12693
15. Pastorello EA, Farioli L, Pravettoni V, et al. Sensitization to the major allergen of Brazil nut is correlated with the clinical expression of allergy. *J Allergy Clin Immunol*. 1998;102(6 Pt 1):1021-1027. doi:10.1016/s0091-6749(98)70341-0
16. Moreno FJ, Clemente A. 2S Albumin storage proteins: What makes them food allergens? *Open Biochem J*. 2008;2:16-28. doi:10.2174/1874091X00802010016

## Performance

### Method Description

Specific IgE from the patient's serum reacts with the allergen of interest, which is covalently coupled to an ImmunoCAP. After washing away nonspecific IgE, enzyme-labeled anti-IgE antibody is added to form a complex. After incubation, unbound anti-IgE is washed away, and the bound complex is then incubated with a developing agent. After stopping the reaction, the fluorescence of the eluate is measured. Fluorescence is proportional to the amount of specific IgE present in the patient's sample (ie, the higher the fluorescence value, the more IgE antibody is present). (Package insert: ImmunoCAP System Specific IgE FEIA. Rev 06/2020)

### PDF Report

No

### Day(s) Performed

Monday through Friday

**Report Available**

Same day/1 to 3 days

**Specimen Retention Time**

14 days

**Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Superior Drive

**Fees & Codes****Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**Test Classification**

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

**CPT Code Information**

86008

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
BRAZX	Brazil Nut Component, IgE, S	64963-2

Result ID	Test Result Name	Result LOINC® Value
IBRAZ	Brazil Nut IgE Ab Interpretation	69048-7
E1BER	Ber e 1 (Brazil Nut), IgE, S	64963-2