

---

## Overview

### Useful For

Evaluating protein S deficiency in patients with a personal or family history suggestive of this hereditary thrombophilia

Confirming a diagnosis of autosomal dominant protein S deficiency with the identification of a known or suspected disease-causing alteration in the *PROS1* gene

Confirming a diagnosis of autosomal recessive severe protein S deficiency with the identification of homozygous or compound heterozygous disease-causing alterations in the *PROS1* gene

Determining the disease-causing alterations within the *PROS1* gene to delineate the underlying molecular defect in a patient with a laboratory diagnosis of protein S deficiency

Prognosis and risk assessment based on genotype-phenotype correlations

Ascertaining the variant status of family members related to an individual with a confirmed *PROS1* variant for the purposes of informing clinical management and genetic counseling

Carrier testing for close family members of an individual with a diagnosis of autosomal recessive severe protein S deficiency

This test is **not intended for** prenatal diagnosis.

### Genetics Test Information

This test utilizes next-generation sequencing to detect single nucleotide and copy number variants in the *PROS1* gene associated with thrombophilia due to protein S deficiency, a rare clotting disorder. See Method Description for additional details.

Identification of a disease-causing variant may assist with diagnosis, prognosis, clinical management, recurrence risk assessment, familial screening, and genetic counseling for thrombophilia due to protein S deficiency.

### Testing Algorithm

The clinical workup for protein S deficiency should begin with measurement of plasma free protein S antigen.

Genetic testing for protein S deficiency is indicated if:

- Free protein S antigen and/or activity is abnormally reduced
- There is a clinical suspicion of hereditary thrombophilia and possible protein S deficiency due to family history or atypical clinical presentation
- Preanalytical variables and acquired causes of protein S deficiency have been excluded (eg, acute thrombosis, surgery, disseminated intravascular coagulation, liver disease, vitamin K deficiency, therapy with vitamin K antagonists such as warfarin, pregnancy, hormonal contraceptives, estrogen therapy, HIV infection, varicella, sickle cell disease, malignancy,

---

nephrotic syndrome)

**Special Instructions**

- [Informed Consent for Genetic Testing](#)
- [Informed Consent for Genetic Testing \(Spanish\)](#)
- [Rare Coagulation Disorder Patient Information](#)

**Method Name**

Sequence Capture and Targeted Next-Generation Sequencing (NGS) followed by Polymerase Chain Reaction (PCR) and Sanger Sequencing

**NY State Available**

Yes

**Specimen****Specimen Type**

Varies

**Ordering Guidance**

This test should only be considered if clinical and family history, initial coagulation screens, and/or initial antigen and activity tests suggest a diagnosis of protein S deficiency (see Testing Algorithm).

This test does not measure protein S activity or antigen levels.

-For assessment of free protein S activity, order S\_FX / Protein S Activity, Plasma.

-For assessment of plasma free protein S antigen, order PSTF / Protein S Antigen, Plasma.

For patients in whom hereditary protein S deficiency is strongly suspected and the plasma free protein S antigen level is normal, consider testing free protein S activity for detecting type II protein S deficiency, which is very rare. Order S\_FX / Protein S Activity, Plasma.

If genetic testing for hereditary blood clotting disorders using a larger panel is desired, a 16-gene comprehensive thrombosis panel is available; order GNTHR / Thrombosis Disorders, Comprehensive Gene Panel, Next-Generation Sequencing, Varies.

Targeted testing for familial variants (also called site-specific or known variants testing) is available for the *PROS1* gene. See FMTT / Familial Variant, Targeted Testing, Varies. To obtain more information about this testing option, call 800-533-1710.

**Necessary Information**

[Rare Coagulation Disorder Patient Information](#) is required. Testing may proceed without the patient information, however, the information aids in providing a more thorough interpretation. Ordering healthcare professionals are

strongly encouraged to fill out the form and send with the specimen.

**Specimen Required**

**Specimen Type:** Whole blood

**Patient Preparation:** A previous bone marrow transplant from an allogenic donor will interfere with testing. For information about testing patients who have received a bone marrow transplant, call 800-533-1710.

**Container/Tube:**

**Preferred:** Lavender top (EDTA)

**Acceptable:** Yellow top (ACD)

**Specimen Volume:** 3 mL

**Collection Instructions:**

1. Invert several times to mix blood.
2. Send whole blood specimen in original tube. **Do not aliquot.**

**Specimen Stability Information:** Ambient (preferred) 4 days/Refrigerated 4 days

**Additional Information:** To ensure minimum volume and concentration of DNA are met, the requested volume must be submitted. Testing may be canceled if DNA requirements are inadequate.

**Forms**

1. [Rare Coagulation Disorder Patient Information \(T824\)](#) is required.
2. **New York Clients-Informed consent is required.** Document on the request form or electronic order that a copy is on file. The following documents are available:
  - [Informed Consent for Genetic Testing \(T576\)](#)
  - [Informed Consent for Genetic Testing \(Spanish\) \(T826\)](#)
3. If not ordering electronically, complete, print, and send an [Coagulation Test Request \(T753\)](#) with the specimen.

**Specimen Minimum Volume**

1 mL

**Reject Due To**

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

**Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Varies	Varies		

**Clinical & Interpretive****Clinical Information**

Hereditary protein S deficiency is a rare inherited clotting disorder (thrombophilia) associated with germline variants in the *PROS1* gene. Protein S is an important component of the body's natural anticoagulant system. A deficiency results in an unchecked clotting cascade and increased risk of thromboembolism.(1)

Mild protein S deficiency is caused by heterozygous variants in one copy of *PROS1* and is inherited in an autosomal

---

dominant manner with incomplete penetrance. Individuals with mild protein S deficiency have a predisposition to developing venous thromboembolism and often experience deep vein thrombosis and/or pulmonary embolism. Affected women have an increased risk for pregnancy loss and complications. The estimated prevalence is 0.16% to 0.21%.(1-4)

Severe protein S deficiency is caused by homozygous or compound heterozygous variants in *PROS1* and is inherited in an autosomal recessive manner. It is quite rare and presents with neonatal purpura fulminans and disseminated intravascular coagulation.(1-3)

Causes of acquired (nongenetic) protein S deficiency should be excluded prior to genetic testing, including vitamin K deficiency, oral anticoagulant therapy, presence of liver disease, intravascular coagulation and fibrinolysis/disseminated intravascular coagulation, thrombotic thrombocytopenia purpura, pregnancy, and estrogen therapy. As an acute-phase reactant, plasma C4b-binding protein levels increase with acute illness and may cause acquired free protein S deficiency.(2)

The British Society for Haematology provides guidelines regarding diagnosis, management, and laboratory testing for individuals with hereditary thrombophilias including protein S deficiency.(5)

### Reference Values

An interpretive report will be provided.

### Interpretation

All detected variants are evaluated according to American College of Medical Genetics and Genomics recommendations.(6) Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.

### Cautions

Clinical Correlations:

Test results should be interpreted in the context of clinical findings, family history, and other laboratory data. Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

If testing was performed because of a clinically significant family history, it is often useful to first test an affected family member. Detection of a reportable variant in an affected family member would allow for more informative testing of at-risk individuals.

To discuss the availability of additional testing options or for assistance in the interpretation of these results, contact the Mayo Clinic Laboratories genetic counselors at 800-533-1710.

Technical Limitations:

Next-generation sequencing may not detect all types of genomic variants. In rare cases, false-negative or false-positive results may occur. The depth of coverage may be variable for some target regions; assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be considered.

---

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. Confirmation of select reportable variants will be performed by alternate methodologies based on internal laboratory criteria.

This test is validated to detect 95% of deletions up to 75 base pairs (bp) and insertions up to 47 bp. Deletions-insertions (delins) of 40 or more bp, including mobile element insertions, may be less reliably detected than smaller delins.

This analysis targets single and multi-exon deletions/duplications; however, in some instances, single exon resolution cannot be achieved due to isolated reduction in sequence coverage or inherent genomic complexity. Balanced structural rearrangements (such as translocations and inversions) may not be detected.

Deletion/duplication events that extend past the gene tested may occur. In these instances, only the gene included in the ordered test is provided on the report and interpreted, and genomic breakpoints are reported if they are confirmed. Genes not on this ordered test are typically not reported or interpreted for haploinsufficiency/triplosensitivity. CMACB / Chromosomal Microarray, Congenital, Blood; WESPR / Panel to Whole Exome Sequencing Reflex Test, Varies; or WGSDX / Whole Genome Sequencing for Hereditary Disorders, Varies is recommended for a full interpretation of deletions/duplications predicted to extend past the tested gene.

This test is not designed to detect low levels of mosaicism or to differentiate between somatic mutations and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

For detailed information regarding gene specific performance and technical limitations, see Method Description or contact a laboratory genetic counselor.

If the patient has had an allogeneic hematopoietic stem cell transplant or a recent blood transfusion, results may be inaccurate due to the presence of donor DNA. Call Mayo Clinic Laboratories for instructions for testing patients who have received a bone marrow transplant.

#### Reclassification of Variants:

Currently, it is not standard practice for the laboratory to systematically review previously classified variants on a regular basis. The laboratory encourages healthcare professionals to contact the laboratory at any time to learn how the classification of a particular variant may have changed over time. Due to broadening genetic knowledge, it is possible that the laboratory may discover new information of relevance to the patient. Should that occur, the laboratory may issue an amended report.

#### Variant Evaluation:

Evaluation and categorization of variants is performed using published American College of Medical Genetics and Genomics and the Association for Molecular Pathology recommendations as a guideline.<sup>(6)</sup> Other gene-specific guidelines may also be considered. Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance. Variants classified as benign or likely benign are not reported.

---

Multiple in silico evaluation tools may be used to assist in the interpretation of these results. The accuracy of predictions made by in silico evaluation tools is highly dependent upon the data available for a given gene, and periodic updates to these tools may cause predictions to change over time. Results from in silico evaluation tools should be interpreted with caution and professional clinical judgment.

Rarely, incidental or secondary findings may implicate another predisposition or presence of active disease. These findings will be carefully reviewed to determine whether they will be reported.

### Clinical Reference

1. Gupta A, Tun AM, Gupta K. Protein S Deficiency. In: StatPearls [Internet]. StatPearls Publishing; Updated December 5, 2022. Accessed March 3, 2026. Available at [www.ncbi.nlm.nih.gov/books/NBK544344/](http://www.ncbi.nlm.nih.gov/books/NBK544344/)
2. ten Kate MK, van der Meer J. Protein S deficiency: a clinical perspective. *Haemophilia*. 2008;14(6):1222-1228
3. Garcia de Frutos P, Fuentes-Prior P, Hurtado B, Sala N. Molecular basis of protein S deficiency. *Thromb Haemost*. 2007;98(3):543-556
4. Beauchamp NJ, Dykes AC, Parikh N, Campbell Tait R, Daly ME. The prevalence of, and molecular defects underlying, inherited protein S deficiency in the general population. *Br J Haematol*. 2004;125(5):647-654
5. Arachchilage DJ, Mackillop L, Chandratheva A, Motawani J, MacCallum P, Laffan M. Thrombophilia testing: A British Society for Haematology guideline. *Br J Haematol*. 2022;198(3):443-458
6. Richards S, Aziz N, Bale S, et al. ACMG Laboratory Quality Assurance Committee: Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-424

### Performance

#### Method Description

Next-generation sequencing (NGS) and/or Sanger sequencing are performed to test for the presence of variants in coding regions and intron/exon boundaries of the *PROS1* gene, as well as some other regions that have known disease-causing variants. The human genome reference GRCh37/hg19 build was used for sequence read alignment. At least 99% of the bases are covered at a read depth over 30X. Sensitivity is estimated at above 99% for single nucleotide variants, above 94% for deletions-insertions (delins) less than 40 base pairs (bp), and above 95% for deletions up to 75 bp, and insertions up to 47 bp. NGS and/or a polymerase chain reaction-based quantitative method is performed to test for the presence of deletions and duplications in the *PROS1* gene.

There may be regions of the *PROS1* gene that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. (Unpublished Mayo method)

The reference transcript for *PROS1* is NM\_000313.4. Reference transcript numbers may be updated due to transcript re-versioning. Always refer to the final patient report for gene transcript information referenced at the time of testing. Confirmation of select reportable variants may be performed by alternate methodologies based on internal laboratory criteria.

## PDF Report

Supplemental

## Day(s) Performed

Varies

## Report Available

28 to 42 days

## Specimen Retention Time

Whole blood: 2 weeks (if available); Extracted DNA: 3 months

## Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

### Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

### CPT Code Information

81479

### LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
GNPRS	PROS1 Gene, Full Gene NGS	92994-3

Result ID	Test Result Name	Result LOINC® Value
619188	Test Description	62364-5
619189	Specimen	31208-2
619190	Source	31208-2
619191	Result Summary	50397-9
619192	Result	82939-0
619193	Interpretation	59465-5
619194	Additional Results	82939-0
619195	Resources	99622-3

## Test Definition: GNPRS

Protein S Deficiency, PROS1 Gene,  
Next-Generation Sequencing, Varies

619196	Additional Information	48767-8
619197	Method	85069-3
619198	Genes Analyzed	82939-0
619199	Disclaimer	62364-5
619200	Released By	18771-6