

Wilson Disease, ATP7B Full Gene Sequencing with Deletion/Duplication, Varies

Overview

Useful For

Confirming the diagnosis of Wilson disease

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
CULFB	Fibroblast Culture for	Yes	No
	Genetic Test		

Genetics Test Information

This test utilizes next-generation sequencing to detect single nucleotide, deletion-insertion, and copy number variants in the *ATP7B* gene associated with Wilson disease. See Method Description for additional details.

Identification of a disease-causing variant may assist with diagnosis, prognosis, clinical management, recurrence risk assessment, familial screening, and genetic counseling for Wilson disease.

Testing Algorithm

For skin biopsy or cultured fibroblast specimens, fibroblast culture testing will be performed at an additional charge. If viable cells are not obtained, the client will be notified.

Special Instructions

- Molecular Genetics: Biochemical Disorders Patient Information
- Informed Consent for Genetic Testing
- Wilson Disease Testing Algorithm
- Informed Consent for Genetic Testing (Spanish)

Method Name

Sequence Capture and Targeted Next-Generation Sequencing (NGS) followed by Polymerase Chain Reaction (PCR) and Sanger Sequencing.

NY State Available

Yes

Specimen

Specimen Type

Varies



Wilson Disease, ATP7B Full Gene Sequencing with Deletion/Duplication, Varies

Ordering Guidance

Testing for the *ATP7B* gene as a part of a customized panel is available. For more information see CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies.

Targeted testing (also called site-specific or known variants testing) is available for variants identified in the *ATP7B* gene. See FMTT / Familial Variant, Targeted Testing, Varies. To obtain more information about this testing option, call 800-533-1710.

Shipping Instructions

Specimen preferred to arrive within 96 hours of collection.

Specimen Required

Patient Preparation: A previous bone marrow transplant from an allogenic donor will interfere with testing. For instructions for testing patients who have received a bone marrow transplant, call 800-533-1710.

Submit only 1 of the following specimens:

Specimen Type: Whole blood

Container/Tube: Lavender top (EDTA) or yellow top (ACD)

Specimen Volume: 3 mL **Collection Instructions:**

1. Invert several times to mix blood.

2. Send whole blood specimen in original tube. **Do not aliquot. Specimen Stability Information:** Ambient (preferred)/Refrigerated

Specimen Type: Blood spot

Container/Tube:

Preferred: Collection card (Whatman Protein Saver 903 Paper)

Acceptable: PerkinElmer 226 (formerly Ahlstrom 226) filter paper, or Blood Spot Collection Card (T493)

Specimen Volume: 5 Blood spots

Collection Instructions:

- 1. An alternative blood collection option for a patient older than 1 year is a fingerstick. For detailed instructions, see How to Collect Dried Blood Spot Samples.
- 2. Let blood dry on the filter paper at ambient temperature in a horizontal position for a minimum of 3 hours.
- 3. Do not expose specimen to heat or direct sunlight.
- 4. Do not stack wet specimens.
- 5. Keep specimen dry

Specimen Stability Information: Ambient (preferred)/Refrigerated

Additional Information:

- 1. Due to lower concentration of DNA yielded from blood spot, it is possible that additional specimen may be required to complete testing.
- 2. For collection instructions, see <u>Blood Spot Collection Instructions</u>
- 3. For collection instructions in Spanish, see Blood Spot Collection Card-Spanish Instructions (T777)
- 4. For collection instructions in Chinese, see Blood Spot Collection Card-Chinese Instructions (T800)



Wilson Disease, ATP7B Full Gene Sequencing with Deletion/Duplication, Varies

Specimen Type: Saliva

Patient Preparation: Patient should not eat, drink, smoke, or chew gum 30 minutes prior to collection.

Supplies:

DNA Saliva Kit High Yield (T1007) Saliva Swab Collection Kit (T786)

Container/Tube:

Preferred: High-yield DNA saliva kit

Acceptable: Saliva swab

Specimen Volume: 1 Tube if using T1007 or 2 swabs if using T786 **Collection Instructions:** Collect and send specimen per kit instructions.

Specimen Stability Information: Ambient (preferred) 30 days/Refrigerated 30 days

Additional Information: Saliva specimens are acceptable but not recommended. Due to lower quantity/quality of DNA yielded from saliva, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.

Specimen Type: Skin biopsy

Supplies: Fibroblast Biopsy Transport Media (T115)

Container/Tube: Sterile container with any standard cell culture media (eg, minimal essential media, RPMI 1640). The

solution should be supplemented with 1% penicillin and streptomycin.

Specimen Volume: 4-mm punch

Specimen Stability Information: Refrigerated (preferred)/Ambient

Additional Information: A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or

Molecular Testing. An additional 3 to 4 weeks is required to culture fibroblasts before genetic testing can occur.

Specimen Type: Cultured fibroblast

Container/Tube: T-25 flask Specimen Volume: 2 Flasks

Collection Instructions: Submit confluent cultured fibroblast cells from a skin biopsy from another laboratory. Cultured

cells from a prenatal specimen will not be accepted.

Specimen Stability Information: Ambient (preferred)/Refrigerated (<24 hours)

Additional Information: A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or

Molecular Testing. An additional 3 to 4 weeks is required to culture fibroblasts before genetic testing can occur.

Forms

1. New York Clients-Informed consent is required.

Document on the request form or electronic order that a copy is on file. The following documents are available:

- -Informed Consent for Genetic Testing (T576)
- -Informed Consent for Genetic Testing (Spanish) (T826)
- 2. Molecular Genetics: Biochemical Disorders Patient Information (T527)
- 3. If not ordering electronically, complete, print, and send 1 of the following forms with the specimen:
- -Gastroenterology and Hepatology Test Request (T728)
- -Biochemical Genetics Test Request (T798)



Wilson Disease, ATP7B Full Gene Sequencing with Deletion/Duplication, Varies

Specimen Minimum Volume

Blood: 1 mL; Blood spots: 2 spots; Skin biopsy, cultured fibroblasts, or saliva: See Specimen Required

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Varies	Varies		

Clinical & Interpretive

Clinical Information

Wilson disease (WD) is an autosomal recessive disorder that results from the body's inability to excrete excess copper. Typically, the liver releases excess copper into the bile. Individuals with WD lack the necessary enzyme that facilitates clearance of copper from the liver to bile. As a result, copper accumulates first in the liver and gradually in other organs. The brain, kidneys, bones, and corneas can also be affected. WD affects approximately 1 in 30,000 people worldwide, with a carrier frequency of approximately 1 in 90 individuals.

The primary clinical manifestations of WD are hepatic and neurologic. Hepatic disease can be quite variable, ranging from hepatomegaly or other nonspecific symptoms that mimic viral hepatitis to severe liver damage, such as cirrhosis. Neurologic symptoms of WD can include poor fine-motor coordination, ataxia, and dysphagia. Psychiatric manifestations are reported in approximately 20% of individuals with WD. A characteristic ophthalmologic finding is the Kayser-Fleischer ring. Individuals with WD typically begin to show symptoms of liver dysfunction or neurologic disease in the first or second decade of life. If not treated, WD can cause liver failure, severe brain damage, and even death.

A variety of laboratory tests are recommended in the initial evaluation for WD. In approximately 95% of cases, serum ceruloplasmin is below normal. Additionally, patients with WD show decreased copper in serum, increased copper in urine, and significantly elevated copper on liver biopsy. While liver biopsy is not recommended as a first-tier screening test for WD, it can be useful to help interpret discrepant biochemical or molecular results. Analyte screening tests should be considered prior to molecular analysis. WD is caused by disease-causing variants in the *ATP7B* gene. More than 300 disease-causing variants have been identified in the *ATP7B* gene. Most disease-causing variants are family-specific, with the exception of the H1069Q variant, which accounts for greater than 50% of identified disease alleles in the Northern European population.

Reference Values

An interpretive report will be provided.

Interpretation

All detected variants are evaluated according to American College of Medical Genetics and Genomics recommendations.(1) Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.



Wilson Disease, ATP7B Full Gene Sequencing with Deletion/Duplication, Varies

Cautions

Clinical Correlations:

Test results should be interpreted in the context of clinical findings, family history, and other laboratory data. Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

If testing was performed because of clinically significant family history, it is often useful to first test an affected family member. Detection of a reportable variant in an affected family member would allow for more informative testing of at-risk individuals.

To discuss the availability of additional testing options or for assistance in the interpretation of these results, contact the Mayo Clinic Laboratories genetic counselors at 800-533-1710.

Technical Limitations:

Next-generation sequencing may not detect all types of genomic variants. In rare cases, false-negative or false-positive results may occur. The depth of coverage may be variable for some target regions; assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be considered.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. Confirmation of select reportable variants will be performed by alternate methodologies based on internal laboratory criteria.

This test is validated to detect 95% of deletions up to 75 base pairs (bp) and insertions up to 47 bp. Deletions-insertions (delins) of 40 or more bp, including mobile element insertions, may be less reliably detected than smaller delins.

Deletion/Duplication Analysis:

This analysis targets single and multi-exon deletions/duplications; however, in some instances, single exon resolution cannot be achieved due to isolated reduction in sequence coverage or inherent genomic complexity. Balanced structural rearrangements (such as translocations and inversions) may not be detected.

This test is not designed to detect low levels of mosaicism or to differentiate between somatic and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

For detailed information regarding gene-specific performance and technical limitations, see Method Description or contact a laboratory genetic counselor.

If the patient has had an allogeneic hematopoietic stem cell transplant or a recent blood transfusion, results may be inaccurate due to the presence of donor DNA. Call Mayo Clinic Laboratories for instructions for testing patients who have received a bone marrow transplant.



Wilson Disease, ATP7B Full Gene Sequencing with Deletion/Duplication, Varies

Reclassification of Variants:

Currently, it is not standard practice for the laboratory to systematically review previously classified variants on a regular basis. The laboratory encourages healthcare providers to contact the laboratory at any time to learn how the classification of a particular variant may have changed over time. Due to broadening genetic knowledge, it is possible that the laboratory may discover new information of relevance to the patient. Should that occur, the laboratory may issue an amended report.

Variant Evaluation:

Evaluation and categorization of variants are performed using published American College of Medical Genetics and Genomics and the Association for Molecular Pathology recommendations as a guideline. (1) Other gene-specific guidelines may also be considered. Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance. Variants classified as benign or likely benign are not reported.

Multiple in silico evaluation tools may be used to assist in the interpretation of these results. The accuracy of predictions made by in silico evaluation tools is highly dependent upon the data available for a given gene, and periodic updates to these tools may cause predictions to change over time. Results from in silico evaluation tools should be interpreted with caution and professional clinical judgment.

Rarely, incidental or secondary findings may implicate another predisposition or presence of active disease. Incidental findings may include, but are not limited to, results related to the sex chromosomes. These findings will be carefully reviewed to determine whether they will be reported.

Clinical Reference

- 1. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. Genet Med. 2015;17(5):405-424
- 2. European Association for Study of Liver. EASL Clinical Practice Guidelines. Wilson's disease. J Hepatol. 2012;56(3):671-685
- 3. Roberts EA, Schilsky ML, American Association for Study of Liver Diseases (AASLD). Diagnosis and treatment of Wilson disease: an update. Hepatology. 200847(6):2089-2111
- 4. Mak CM, Lam CW. Diagnosis of Wilson's disease: a comprehensive review. Crit Rev Clin Lab Sci. 2008;45(3):263-290
- 5. Bandmann O, Weiss KH, Kaler SG. Wilson's disease and other neurological copper disorders. Lancet Neurol. 2015;14(1):103-113

Performance

Method Description

Next-generation sequencing (NGS) and Sanger sequencing are performed to test for the presence of variants in coding regions and intron/exon boundaries of *ATP7B* gene, as well as some other regions that have known disease-causing variants. The human genome reference GRCh37/hg19 build was used for sequence read alignment. At least 99% of the bases are covered at a read depth over 30X. Sensitivity is estimated to be over 99% for single nucleotide variants, over



Wilson Disease, ATP7B Full Gene Sequencing with Deletion/Duplication, Varies

94% for deletions/insertions (delins) less than 40 base pairs (bp), over 95% for deletions up to 75 bp and insertions up to 47 bp. NGS and/or a polymerase chain reaction-based quantitative method is performed to test for the presence of deletions and duplications in the *ATP7B* gene.

There may be regions of this gene that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. (Unpublished Mayo method)

The reference transcript for the *ATP7B* gene is NM_000053.4. Reference transcript numbers may be updated due to transcript re-versioning. Always refer to the final patient report for gene transcript information referenced at the time of testing. Confirmation of select reportable variants may be performed by alternate methodologies based on internal laboratory criteria.

The following additional noncoding variants are being analyzed by this test: c.-676A>G, c.-460G>A, c.-447C>T, c.-442G>A, c.-436_-422del15, c.-388C>T, c.-210A>T, c.-133A>C, c.-128A>C, c.-123C>A, and c.-128_-124delAGCCG.

PDF Report

Supplemental

Day(s) Performed

Varies

Report Available

14 to 21 days

Specimen Retention Time

Whole blood: 2 weeks (if available); Extracted DNA: 3 months; Blood spots, saliva, cultured fibroblasts, skin biopsy: 1 month

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.



Wilson Disease, ATP7B Full Gene Sequencing with Deletion/Duplication, Varies

CPT Code Information

81406

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
WNDZ	ATP7B Full Gene Analysis	95781-1

Result ID	Test Result Name	Result LOINC® Value
619410	Test Description	62364-5
619411	Specimen	31208-2
619412	Source	31208-2
619413	Result Summary	50397-9
619414	Result	82939-0
619415	Interpretation	69047-9
619416	Additional Results	82939-0
619417	Resources	99622-3
619418	Additional Information	48767-8
619419	Method	85069-3
619420	Genes Analyzed	48018-6
619421	Disclaimer	62364-5
619422	Released By	18771-6