

**Overview****Useful For**

Lymphoma classification

**Reflex Tests**

Test Id	Reporting Name	Available Separately	Always Performed
IHTOI	IHC Initial, Tech Only	No	No
IHTOA	IHC Additional, Tech Only	No	No

**Testing Algorithm**

For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

**Method Name**

Immunohistochemistry (IHC)

**NY State Available**

Yes

**Specimen****Specimen Type**

TECHONLY

**Ordering Guidance**

This test includes only technical performance of the stain (no pathologist interpretation is performed). If diagnostic consultation by a pathologist is required order PATHC / Pathology Consultation.

**Shipping Instructions**

Attach the green pathology address label and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

**Specimen Required**

**Specimen Type:** Tissue

**Supplies:** Immunostain Technical Only Envelope (T693)

**Container/Tube:** Immunostain Technical Only Envelope

**Preferred:**

-Formalin-fixed, paraffin-embedded tissue block

OR

-2 Unstained, positively charged glass slides (25- x 75- x 1-mm) per test ordered; sections 4-microns thick

**Acceptable:** None

### Digital Image Access

1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: <https://news.mayocliniclabs.com/ihc-stains/>
2. Clients ordering stains using a manual requisition form will not have access to digital images.
3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: <https://news.mayocliniclabs.com/ihc-stains/#FAQ>

### Forms

If not ordering electronically, complete, print, and send a [Immunohistochemical \(IHC\)/In Situ Hybridization \(ISH\) Stains Request](#) (T763) with the specimen.

### Reject Due To

Wet/frozen tissue Cytology smears Nonformalin fixed tissue Nonparaffin embedded tissue Noncharged slides ProbeOn slides Snowcoat slides	Reject
---	--------

### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

### Clinical & Interpretive

#### Clinical Information

Myocyte enhancer factor 2 (MEF2B) is a marker of normal germinal center (GC) B cells and lymphomas derived from GC

B cells. MEF2B expression is found in follicular lymphoma, Burkitt lymphoma, and nodular lymphocyte predominant Hodgkin lymphoma, as well as in most cases of mantle cell lymphoma and diffuse large cell B-cell lymphoma.

**Interpretation**

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

**Cautions**

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

The charge of glass slides can be affected by environmental factors and subsequently may alter slide staining. Sending unsuitable glass slides can result in inconsistent staining due to poor slide surface chemistry.

Best practices for storage of positively charged slides:

- Minimize time slides are stored after being unpackaged
- Limit exposure to high humidity and heat
- Minimize exposure to plastics

**Clinical Reference**

1. Krenacs D, Borbenyi Z, Bedekovics J, Mehes G, Bagdi E, Krenacs L: Pattern of MEF2B expression in lymphoid tissues and in malignant lymphomas. *Virchows Arch.* 2015 Sep;467(3):345-355
2. Moore EM, Swerdlow SH, Gibson SE: Comparison of myocyte enhancer factor 2B versus other germinal center-associated antigens in the differential diagnosis of B-cell non-Hodgkin lymphomas. *Am J Surg Pathol.* 2018 Mar;42(3):342-350
3. El Jamal SM, Grada Z, El Dinali MH, et al: MEF2B is a member of the BCL6 gene transcriptional complex and induces its expression in diffuse large B-cell lymphoma of the germinal center B-cell-like type. *Lab Invest.* 2019 Apr;99(4):539-550
4. Salama A, Marcellino BK, Saad AG, et al: Correlation analysis between the expression of MEF2B, and germinal center and nongerminal center markers in diffuse large B-cell lymphoma. *Appl Immunohistochem Mol Morphol.* 2020 Aug;28(7):e63-e64

**Performance****Method Description**

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

**PDF Report**

No

**Day(s) Performed**

Monday through Friday

**Report Available**

1 to 3 days

**Specimen Retention Time**

Until staining is complete

**Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

**Fees & Codes****Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**Test Classification**

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

**CPT Code Information**

88342-TC, primary

88341-TC, if additional IHC

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
MEF2B	MEF2B IHC, Tech Only	No LOINC Needed

Result ID	Test Result Name	Result LOINC® Value
619495	MEF2B IHC, Tech Only	No LOINC Needed