

Mycobacterium tuberculosis Complex, Molecular Detection and Rifampin Resistance, PCR, Sputum

#### Overview

#### **Useful For**

Rapid detection of *Mycobacterium tuberculosis* DNA from respiratory specimens for the diagnosis of pulmonary tuberculosis

Presumptive detection of rifampin resistance based on the presence of resistance-associated mutations

#### **Method Name**

Real-Time Polymerase Chain Reaction (PCR)

#### **NY State Available**

Yes

# Specimen

## **Specimen Type**

Sputum

## **Additional Testing Requirements**

Mycobacterial culture is required for epidemiological strain typing and growth-based phenotypic antimicrobial susceptibility testing, including definitive rifampin results as well as results for other antimicrobials. If your facility is unable to perform a mycobacterial culture, order CTB / Mycobacteria and *Nocardia* Culture, Varies concurrently with this test.

### Shipping Instructions

Specimen must arrive within 7 days of collection; if received greater than 7 days from collection, the specimen will be rejected.

# **Necessary Information**

Specimen source is required.

#### Specimen Required

The high sensitivity of amplification by polymerase chain reaction requires the specimen to be processed in an environment in which contamination of the specimen by *Mycobacterium tuberculosis* DNA is unlikely.

**Specimen Type:** Sputum (undigested) **Container/Tube:** Sterile container

Specimen Volume: 3 mL



Mycobacterium tuberculosis Complex, Molecular Detection and Rifampin Resistance, PCR, Sputum

Specimen Stability Information: Refrigerated (preferred) 7 days/Ambient 72 hours

#### **Additional Information:**

- 1. If a single specimen is being shared between mycobacterial culture, acid-fast smear, and/or *M tuberculosis* PCR, a minimum volume of 3 mL for the respiratory specimen is required. Specimen volumes less than indicated may decrease sensitivity of testing.
- 2. If insufficient volume is submitted, testing will be canceled.

Specimen Type: N-acetyl-L-cysteine/sodium hydroxide (NALC/NaOH)-digested sputum

Container/Tube: Sterile container

Specimen Volume: 3 mL Collection Instructions:

1. Submit digested specimen treated with NALC/NaOH.

2. Clearly indicate on container and order form that specimen is a digested specimen.

Specimen Stability Information: Refrigerated 7 days

### **Additional Information:**

- 1. If a single specimen is being shared between mycobacterial culture, acid-fast smear, and/or *M tuberculosis* PCR, a minimum volume of 3 mL for the respiratory specimen is required. Specimen volumes less than indicated may decrease sensitivity of testing.
- 2. If insufficient volume is submitted, testing will be canceled.

#### **Forms**

If not ordering electronically, complete, print, and send a Microbiology Test Request (T244) with the specimen.

## **Specimen Minimum Volume**

1.5 mL

### Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

#### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Sputum	Refrigerated (preferred)	7 days	
	Ambient	72 hours	

# Clinical & Interpretive

## **Clinical Information**

Mycobacterium tuberculosis is a highly transmissible bacterial pathogen and is the causative agent of tuberculosis, a disease causing significant worldwide morbidity and mortality. Each year, M tuberculosis accounts for 1.6 million deaths and is responsible for 10.6 million newly diagnosed cases of tuberculosis worldwide. M tuberculosis is spread from person to person via respiratory transmission and has the potential to become resistant to many of the antibiotics



Mycobacterium tuberculosis Complex, Molecular Detection and Rifampin Resistance, PCR, Sputum

currently used if not treated appropriately. Therefore, rapid and accurate detection of *M tuberculosis* in patient specimens is of paramount clinical and public health importance.

Conventional culture methods can generally detect *M tuberculosis* in 2 to 3 weeks, although up to 6 weeks of incubation may be required in some instances. This qualitative molecular assay utilizes polymerase chain reaction-based nucleic acid amplification for the direct detection of *M tuberculosis* DNA within respiratory specimens without relying on culture growth, leading to more rapid diagnosis and appropriate patient care. This assay also detects the presence of mutations in the *rpoB* gene that have been documented to confer more than 95% of cases of rifampin resistance.

#### **Reference Values**

Negative

### Interpretation

A positive result indicates the presence of Mycobacterium tuberculosis complex DNA.

A negative result indicates the absence of detectable M tuberculosis complex DNA.

Presumptive rifampin resistance mediated through mutations within the resistance determining region of the *rpoB* gene will be reported when detected.

One to 2 negative polymerase chain reaction results in conjunction with 1 to 2 negative acid-fast smears may provide evidence supporting the removal of a patient from airborne isolation. Consult your local Infection Prevention and Control for guidance.

#### Cautions

This test should always be performed in conjunction with mycobacterial culture, which is required for epidemiological strain typing and growth-based phenotypic antimicrobial susceptibility testing, including definitive rifampin results as well as results for other antimicrobials. If your facility is unable to perform mycobacterial culture, CTB / Mycobacteria and *Nocardia* Culture, Varies should be ordered.

Per current Centers for Disease Control and Prevention recommendations, rifampin resistance results should be considered as preliminary pending definitive confirmation with gene sequencing or growth-based phenotypic antimicrobial susceptibility testing.

This polymerase chain reaction-based molecular assay detects *Mycobacterium tuberculosis* nucleic acid and, therefore, does not distinguish between viable, disease-related organisms and nucleic acid persisting from prior infection. Test results should be correlated with patient symptoms and clinical presentation before a definitive diagnosis is made.

A negative result does not rule-out infection with *M tuberculosis* or active disease because the organism may be present at levels below the limit of detection for this assay.

# **Clinical Reference**

1. World Health Organization. Global Tuberculosis Report 2022. WHO; 2022 Available at www.who.int/publications/i/item/9789240061729



Mycobacterium tuberculosis Complex, Molecular Detection and Rifampin Resistance, PCR, Sputum

- 2. Centers for Disease Control and Prevention (CDC). Availability of an assay for detecting *Mycobacterium tuberculosis*, including rifampin-resistant strains, and considerations for its use United States, 2013. MMWR Morb Mortal Wkly Rep. 2013 Oct 18;62(41):821-827
- 3. Boehme CC, Nabeta P, Hillemann D, et al. Rapid molecular detection of tuberculosis and rifampin resistance. N Engl J Med. 2010;363(11):1005-1015
- 4. US Food and Drug Administration. New data shows test can help physicians remove patients with suspected TB from isolation earlier. Press Release. 2015. Available at

www.tbonline.info/posts/2015/2/12/fda-new-data-shows-test-can-help-physicians-remove/

#### **Performance**

## **Method Description**

The Cepheid Xpert MTB/RIF assay utilizes sputum or decontaminated and pelleted sputum sediment. Specimens are inoculated directly into single disposable test cartridges, which contain both DNA extraction and target amplification material. The assay amplifies a 192 base-pair segment of the *rpoB* gene for *Mycobacterium tuberculosis* complex identification as well as rifampin resistance profiling. The Cepheid Xpert MTB/RIF assay is a closed polymerase chain reaction (PCR) system that greatly reduces the potential for false-positive results due to specimen cross-contamination as compared with traditional open-system PCR methods.(Package insert: Xpert MTB/RIF. Cepheid; GXMTB/RIF-US-10, rev D, 03/2016)

## **PDF Report**

No

### Day(s) Performed

Monday through Sunday

## Report Available

Same day/1 day

# **Specimen Retention Time**

7 days

## **Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

#### Fees & Codes

#### **Fees**

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.



Mycobacterium tuberculosis Complex, Molecular Detection and Rifampin Resistance, PCR, Sputum

• Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

#### **Test Classification**

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

# **CPT Code Information**

87564

## **LOINC®** Information

Test ID	Test Order Name	Order LOINC® Value
MTBXS	M tuberculosis/ RIF PCR GeneXpert	89371-9

Result ID	Test Result Name	Result LOINC® Value
SRCPX	Specimen Source	31208-2
MTBRX	MTB Complex Result	88874-3
RIFRX	MTB Rifampin Resistance Result	89372-7