

## Overview

### Useful For

Identifying specific mutations within the *SDHA*, *SDHB*, *SDHC*, and *SDHD* genes to assist in tumor diagnosis/classification, including pheochromocytoma/paraganglioma, renal cell carcinoma, and pituitary adenoma

### Genetics Test Information

This test uses targeted next-generation sequencing to evaluate for somatic mutations within the *SDHA*, *SDHB*, *SDHC*, and *SDHD* genes. See [Targeted Genes and Methodology Details for SDH Genes Mutation Analysis](#) for details regarding the targeted gene regions evaluated by this test.

This test is performed to evaluate for somatic mutations within solid tumor samples. This test **does not assess** for germline alterations within the genes listed.

### Additional Tests

Test Id	Reporting Name	Available Separately	Always Performed
SLIRV	Slide Review in MG	No, (Bill Only)	Yes

### Testing Algorithm

When this test is ordered, slide review will always be performed at an additional charge.

### Special Instructions

- [Tissue Requirements for Solid Tumor Next-Generation Sequencing](#)
- [Targeted Genes and Methodology Details for SDH Genes Mutation Analysis](#)

### Method Name

Sequence Capture Next-Generation Sequencing (NGS)

### NY State Available

Yes

## Specimen

### Specimen Type

Varies

### Ordering Guidance

Multiple oncology (cancer) gene panels are available. For more information see [Hematology, Oncology, and Hereditary Test Selection Guide](#).

**Necessary Information**

**A pathology report** (final or preliminary), at minimum containing the following information, **must accompany specimen for testing to be performed**:

1. Patient name
2. Block number-must be on all blocks, slides, and paperwork (can be handwritten on the paperwork)
3. Tissue collection date
4. Source of the tissue

**Specimen Required**

**This assay requires at least 20% tumor nuclei.**

- Preferred amount of tumor area with sufficient percent tumor nuclei: tissue 216 mm(2)
- Minimum amount of tumor area: tissue 36 mm(2)
- These amounts are cumulative over up to 10 unstained slides and must have adequate percent tumor nuclei.
- Tissue fixation: 10% neutral buffered formalin, not decalcified
- For specimen preparation guidance, see [Tissue Requirement for Solid Tumor Next-Generation Sequencing](#). In this document, the sizes are given as 4 mm x 4mm x 10 slides as preferred: approximate/equivalent to 144 mm(2) and the minimum as 3 mm x 1 mm x 10 slides: approximate/equivalent to 36 mm(2).

**Preferred:** Submit 3, if available, or 2 of the following specimens.

**Acceptable:** Submit **at least one** of the following specimens.

**Specimen Type:** Tissue block

**Collection Instructions:** Submit a formalin-fixed, paraffin-embedded tissue block with acceptable amount of tumor tissue.

**Specimen Type:** Tissue slide

**Slides:** 1 Hematoxylin and eosin-stained and 10 unstained

**Collection Instructions:**

Submit the following slides:

1 Slide stained with hematoxylin and eosin

AND

15 Unstained, nonbaked slides with 5-micron thick sections of the tumor tissue.

**Note:** The total amount of required tumor nuclei can be obtained by scraping up to 10 slides from the same block.

**Additional Information:** Unused unstained slides will not be returned.

**Specimen Type:** Cytology slide (direct smears or ThinPrep)

**Slides:** 1 to 3 Slides

**Collection Instructions:** Submit 1 to 3 slides stained and coverslipped with a total of 5000 nucleated cells (preferred) or at least 3000 nucleated cells (minimum).

**Note:** Glass coverslips are preferred; plastic coverslips are acceptable but will result in longer turnaround times.

**Additional Information:** Cytology slides will not be returned. An image of the slides will be stored per regulatory requirements.

**Forms**

If not ordering electronically, complete, print, and send an [Oncology Test Request](#) (T729) with the specimen.

**Specimen Minimum Volume**

See Specimen Required

**Reject Due To**

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

**Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Varies	Ambient (preferred)		
	Refrigerated		

**Clinical & Interpretive****Clinical Information**

Disease-causing alterations of the succinate dehydrogenase complex genes (including *SDHA*, *SDHB*, *SDHC*, and *SDHD*) have been implicated in multiple tumor types, including pheochromocytoma/paraganglioma, renal cell carcinoma, gastrointestinal stromal tumors, and pituitary adenoma. Germline alterations of the *SDH* genes have been associated with hereditary pheochromocytoma/paraganglioma syndromes. The 5th edition of the World Health Organization classification of tumors recognizes succinate dehydrogenase-deficient renal cell carcinoma as a molecularly defined entity.<sup>(1)</sup> This assay, performed using formalin-fixed paraffin-embedded tissue or cytology material, is therefore helpful in documenting an underlying disease-causing alteration of the *SDH* genes and is diagnostically significant. Note that this assay does not distinguish between germline alterations and somatic mutations or identify epigenetic alterations of interest (such as *SDHC* promoter hypermethylation).

**Reference Values**

An interpretive report will be provided.

**Interpretation**

The interpretation of molecular biomarker analysis includes an overview of the results and the associated diagnostic, prognostic, and therapeutic implications.

**Cautions**

This test cannot differentiate between somatic mutations and germline alterations. Additional testing may be necessary to clarify the significance of results if there is a potential hereditary risk.

DNA variants of uncertain significance may be identified.

A negative result does not rule out the presence of a variant that may be present below the limits of detection of this assay. In a specimen with 20% or more tumor content, the analytical sensitivity of this assay for sequence reportable alterations is 5% mutant allele frequency with a minimum coverage of 500X.

---

Point mutations and small deletion-insertion mutations will be detected in the *SDHA*, *SDHB*, *SDHC*, and *SDHD* genes only. This test may detect single exon deletions but does not detect multi-exon deletions, duplications, epigenetic alterations (such as promoter hypermethylation), or genomic copy number variants.

Variant allele frequency (VAF) is the percentage of sequencing reads supporting a specific variant divided by the total sequencing reads at that position. In somatic testing, VAF should be interpreted in the context of several factors, including, but not limited to, tumor purity/heterogeneity/copy number status (ploidy, gains/losses, loss of heterozygosity) and sequencing artifact/misalignment.(2,3)

Rare genetic alterations (ie, polymorphisms) may be present that could lead to false-negative or false-positive results.

Test results should be interpreted in the context of clinical, tumor sampling, histopathological, and other laboratory data. If results obtained do not match other clinical or laboratory findings, contact the laboratory for discussion. Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

Reliable results are dependent on adequate specimen collection and processing. This test has been validated on cytology slides and formalin-fixed, paraffin-embedded tissues; other types of fixatives are discouraged. Improper treatment of tissues, such as decalcification, may cause polymerase chain reaction failure.

## **Supportive Data**

Performance Characteristics:

The limit of detection for calling a somatic variant (single nucleotide variants [SNV] and deletions-insertions [delins]) is 5% variant allele frequency (VAF) and having at least 500x deduplicated coverage.

Verification studies demonstrated concordance between this test and the reference method for detection of SNV and delins is 98.5% (673/683) and 98.4% (122/124) of variants, respectively. Concordance for the detection of delins was 99.0% (100/101) in variants 1 to 10 base pairs (bp) in size, 93.3% (14/15) in variants 11 to 50 bp in size, and 100% (8/8) in variants over 50 bp in size.

To ensure accuracy, this test will be performed on cases that are estimated by a pathologist to have at least 20% tumor cells.

## **Clinical Reference**

1. WHO Classification of Tumours Editorial Board, eds. Urinary and male genital tumors. 5th ed. World Health Organization; 2022. WHO Classification of Tumours. Vol 8
2. Strom SP. Current practices and guidelines for clinical next-generation sequencing oncology testing. *Cancer Biol Med*. 2016;13(1):3-11. doi:10.28092/j.issn.2095-3941.2016.0004
3. Spurr L, Li M, Alomran N, et al. Systematic pan-cancer analysis of somatic allele frequency. *Sci Rep*. 2018;8(1):7735. Published 2018 May 16. doi:10.1038/s41598-018-25462-0
4. Trpkov K, Hes O, Williamson SR, et al. New developments in existing WHO entities and evolving molecular concepts: The Genitourinary Pathology Society (GUPS) update on renal neoplasia. *Mod Pathol*. 2021;34(7):1392-1424
5. Fuchs TL, Maclean F, Turchini J, et al. Expanding the clinicopathological spectrum of succinate dehydrogenase-deficient renal cell carcinoma with a focus on variant morphologies: a study of 62 new tumors in 59 patients. *Mod Pathol*. 2022;35(6):836-849

6. Gupta S, Swanson AA, Chen YB, et al. Incidence of succinate dehydrogenase and fumarate hydratase-deficient renal cell carcinoma based on immunohistochemical screening with SDHA/SDHB and FH/2SC. *Hum Pathol.* 2019;91:114-122
7. Carlo MI, Hakimi AA, Stewart GD, et al. Familial kidney cancer: Implications of new syndromes and molecular insights. *Eur Urol.* 2019;76(6):754-764
8. Gupta S, Erickson LA. Back to biochemistry: Evaluation for and prognostic significance of SDH mutations in paragangliomas and pheochromocytomas. *Surg Pathol Clin.* 2023;16(1):119-129

## Performance

### Method Description

Next-generation sequencing is performed to evaluate the presence of a mutation in most coding regions of the *SDHA*, *SDHB*, *SDHC*, and *SDHD* genes. See [Targeted Genes and Methodology Details for SDH Genes Mutation Analysis](#) for details regarding the targeted gene regions identified by this test.(Unpublished Mayo method)

A pathology review and macro dissection to enrich tumor cells is performed prior to slide scraping.

### PDF Report

No

### Day(s) Performed

Monday through Friday

### Report Available

12 to 20 days

### Specimen Retention Time

Tissue blocks: Unused portions of blocks will be returned; Tissue slides: Unused slides are stored for at least 5 years; Extracted DNA: 3 months

### Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

### Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA

---

requirements. It has not been cleared or approved by the US Food and Drug Administration.

**CPT Code Information**

88381-Microdissection, manual

81404

81405 x2

81406

81479 (if appropriate for government payers)

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
SDHMP	SDH Genes Mutation Analysis, Tumor	105598-7

Result ID	Test Result Name	Result LOINC® Value
619713	Result	82939-0
619714	Interpretation	69047-9
619715	Additional Information	48767-8
619716	Specimen	31208-2
619717	Tissue ID	80398-1
619718	Method	85069-3
619719	Disclaimer	62364-5
619720	Released By	18771-6