

Overview

Useful For

Providing a comprehensive genetic evaluation for patients with a personal or family history suggestive of an inborn error of immunity causing a hereditary form of severe viral susceptibility

Establishing a diagnosis of hereditary form of viral susceptibility, allowing for appropriate management and surveillance for disease features based on the gene and/or variant involved

Identifying variants within genes known to be associated with a hereditary form of viral susceptibility, allowing for predictive testing of at-risk family members

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
CULFB	Fibroblast Culture for Genetic Test	Yes	No

Genetics Test Information

This test utilizes next-generation sequencing to detect single nucleotide and copy number variants in 30 genes associated with a hereditary form of viral susceptibility: *CARMIL2*, *CD27*, *CD70*, *CIB1*, *CTPS1*, *CXCR4*, *DBR1*, *IFIH1*, *IFNAR1*, *IFNAR2*, *IRF3*, *IRF7*, *IRF9*, *MAGT1*, *POLR3A*, *POLR3C*, *PRKCD*, *RASGRP1*, *SH2D1A*, *STAT1*, *STAT2*, *TLR3*, *TLR7*, *TLR8*, *TMC6*, *TMC8*, *TNFRSF9*, *TRAF3*, *UNC93B1*, and *XIAP*. See [Targeted Genes and Methodology Details for Viral Susceptibility, Defects in Intrinsic and Innate Immunity, Gene Panel](#) and Method Description for additional details.

Identification of a disease-causing variant may assist with diagnosis, prognosis, clinical management, recurrence risk assessment, familial screening, and genetic counseling for a hereditary form of viral susceptibility.

Testing Algorithm

For skin biopsy or cultured fibroblast specimens, fibroblast culture will be performed at an additional charge. If viable cells are not obtained, the client will be notified.

Special Instructions

- [Informed Consent for Genetic Testing](#)
- [Informed Consent for Genetic Testing \(Spanish\)](#)
- [Targeted Genes and Methodology Details for Viral Susceptibility, Defects in Intrinsic and Innate Immunity, Gene Panel](#)
- [Viral Susceptibility, Lymphoproliferation, and Hemophagocytic Lymphohistiocytosis Patient Information](#)

Method Name

Sequence Capture and Targeted Next-Generation Sequencing (NGS)

NY State Available

Yes

Specimen

Specimen Type

Varies

Ordering Guidance

For patients with Epstein-Barr virus (EBV) susceptibility or a heritable predisposition to lymphoproliferative diseases, see EBLPD / Epstein-Barr Virus (EBV) Susceptibility and Lymphoproliferative Disorders Gene Panel, Varies.

Targeted testing for familial variants (also called site-specific or known variants testing) is available for the genes on this panel. See FMTT / Familial Variant, Targeted Testing, Varies. To obtain more information about testing option, call 800-533-1710.

Shipping Instructions

Specimen preferred to arrive within 96 hours of collection.

Specimen Required

Patient Preparation: A previous bone marrow transplant from an allogenic donor will interfere with testing. Call 800-533-1710 for instructions for testing patients who have received a bone marrow transplant.

Submit only 1 of the following specimens:

Specimen Type: Whole blood

Container/Tube:

Preferred: Lavender top (EDTA) or yellow top (ACD)

Acceptable: Any anticoagulant

Specimen Volume: 3 mL

Collection Instructions:

1. Invert several times to mix blood.
2. Send whole blood specimen in original tube. **Do not aliquot.**

Specimen Stability Information: Ambient (preferred) 4 days/Refrigerated

Specimen Type: Skin biopsy

Supplies: Fibroblast Biopsy Transport Media (T115)

Container/Tube: Sterile container with any standard cell culture media (eg, minimal essential media, RPMI 1640). The solution should be supplemented with 1% penicillin and streptomycin.

Specimen Volume: 4-mm punch

Specimen Stability Information: Refrigerated (preferred)/Ambient

Additional Information: A separate culture charge will be assessed under CULFB /Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks is required to culture fibroblasts before genetic testing can occur.

Specimen Type: Cultured fibroblasts

Container/Tube: T-25 flask

Specimen Volume: 2 Flasks

Collection Instructions: Submit confluent cultured fibroblast cells from a skin biopsy from another laboratory. Cultured cells from a prenatal specimen will not be accepted.

Specimen Stability Information: Ambient (preferred)/Refrigerated (<24 hours)

Additional Information: A separate culture charge will be assessed under CULFB /Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks is required to culture fibroblasts before genetic testing can occur.

Forms

- 1. **New York Clients-Informed consent is required.** Document on the request form or electronic order that a copy is on file. The following documents are available:
 - [Informed Consent for Genetic Testing](#) (T576)
 - [Informed Consent for Genetic Testing \(Spanish\)](#) (T826)
- 2. [Molecular Genetics: Congenital Inherited Diseases Patient Information](#) (T521)
- 3. [Viral Susceptibility, Lymphoproliferation, and Hemophagocytic Lymphohistiocytosis Patient Information](#)

Specimen Minimum Volume

Blood: 1 mL; Skin biopsy or cultured fibroblasts: See Specimen Required

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Varies	Varies		

Clinical & Interpretive

Clinical Information

Viral infections are common in otherwise healthy individuals, but they may also present clinically due to a primary (genetic) immunodeficiency (ie, inborn error of immunity: IEI). Alternatively, secondary immunodeficiencies may have a similar presentation but result from immunosuppressive medication or illness, such as HIV infection. IEIs may cause a susceptibility to an entire group of pathogens (bacteria, fungi, or viruses), a subset of pathogens (eg, RNA viruses), or can cause susceptibility specific to a single pathogen (eg, Epstein-Barr virus [EBV], human papillomavirus [HPV]). IEIs may also lead to a more severe presentation, including fatal infection caused by a pathogen that usually causes only a mild or non-fatal disease (eg, influenza). This panel targets IEIs that lead to susceptibility to viruses or to a subset of viruses, severe viral pneumonia, or a specific virus. Examples of infections where this gene panel is useful include EBV,

skin-tropic beta-HPV, influenza, and SARS-CoV-2. IELs that lead to systemic immune deficiencies and susceptibility to a large variety of pathogens (eg, T-cell deficiencies) are not included in this panel.

EBV is the cause of infectious mononucleosis and persists asymptomatically for life in nearly all adults. It is also associated with the development of T- and B-cell lymphomas, nasopharyngeal and gastric carcinomas, and other malignancies. EBV infection in IELs can present with fulminant infectious mononucleosis, hemophagocytosis, B-cell proliferative disease (including lymphoma), and hypogammaglobulinemia.

Beta-HPVs circulate silently in the general population and cause no visible lesions in most people. Genetic susceptibility to beta-HPVs leads to warts, pityriasis-like lesions, epidermodysplasia verruciformis, and increased risk of non-melanoma skin cancers.

Seasonal influenza viruses are common RNA viruses that infect the respiratory tract, causing a benign illness in most infected individuals. Influenza pneumonia or acute respiratory distress syndrome are rare, and the case fatality ratio is less than 1%. Children with severe influenza have been found to carry defects in *IRF7*, *IRF9*, *STAT1*, *STAT2*, and *TLR3*. Similarly, the COVID-19 pandemic has revealed that SARS-CoV-2 infection can lead to asymptomatic infection as well as fatal pneumonia. Genetic studies showed that approximately 2 to 3% of cases of severe life-threatening SARS-CoV-2 infection resulted from IEL, mainly genetic defects in the TLR3- or TLR7-dependent type 1 interferon pathway (eg, *TLR3*, *TLR7*, *IFNAR1/2*, *STAT2*, and *IRF7*), overlapping with that of severe pneumonia susceptibility in influenza infections.

Reference Values

An interpretive report will be provided.

Interpretation

All detected variants are evaluated according to American College of Medical Genetics and Genomics recommendations.(1) Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.

Cautions

Clinical Correlations:

Test results should be interpreted in the context of clinical findings, family history, and other laboratory data.

Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

If testing was performed because of a clinically significant family history, it is often useful to first test an affected family member. Detection of a reportable variant in an affected family member would allow for more informative testing of at-risk individuals.

To discuss the availability of additional testing options or for assistance in the interpretation of these results, contact Mayo Clinic Laboratories genetic counselors at 800-533-1710.

Technical Limitations:

Next-generation sequencing may not detect all types of genomic variants. In rare cases, false-negative or false-positive results may occur. The depth of coverage may be variable for some target regions; assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be

considered.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. Confirmation of select reportable variants will be performed by alternate methodologies based on internal laboratory criteria.

This test is validated to detect 95% of deletions up to 75 base pairs (bp) and insertions up to 47 bp. Deletions-insertions (delins) of 40 or more bp, including mobile element insertions, may be less reliably detected than smaller delins.

Deletion/Duplication Analysis:

This analysis targets single and multi-exon deletions/duplications; however, in some instances, single exon resolution cannot be achieved due to isolated reduction in sequence coverage or inherent genomic complexity. Balanced structural rearrangements (such as translocations and inversions) may not be detected.

This test is not designed to detect low levels of mosaicism or to differentiate between somatic and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

Genes may be added or removed based on updated clinical relevance. Refer to the [Targeted Genes and Methodology Details for Viral Susceptibility, Defects in Intrinsic and Innate Immunity, Gene Panel](#) for the most up to date list of genes included in this test. For detailed information regarding gene-specific performance and technical limitations, see Method Description or contact a laboratory genetic counselor.

[If the patient has had an allogeneic hematopoietic stem cell transplant or a recent non-leukoreduced blood transfusion](#), results may be inaccurate due to the presence of donor DNA. Call Mayo Clinic Laboratories for instructions for testing patients who have received a bone marrow transplant.

Reclassification of Variants:

Currently, it is not standard practice for the laboratory to systematically review previously classified variants on a regular basis. The laboratory encourages health care providers to contact the laboratory at any time to learn how the classification of a particular variant may have changed over time. Due to broadening genetic knowledge, it is possible that the laboratory may discover new information of relevance to the patient. Should that occur, the laboratory may issue an amended report.

Variant Evaluation:

Evaluation and categorization of variants are performed using published American College of Medical Genetics and Genomics and the Association for Molecular Pathology recommendations as a guideline.⁽¹⁾ Other gene-specific guidelines may also be considered. Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance. Variants classified as benign or likely benign are not reported.

Multiple in silico evaluation tools may be used to assist in the interpretation of these results. The accuracy of predictions made by in silico evaluation tools is highly dependent upon the data available for a given gene, and periodic updates to these tools may cause predictions to change over time. Results from in silico evaluation tools should be interpreted with

caution and professional clinical judgment.

Rarely, incidental or secondary findings may implicate another predisposition or presence of active disease. These findings will be carefully reviewed to determine whether they will be reported.

Clinical Reference

1. Richards S, Aziz N, Bale S, et al; ACMG Laboratory Quality Assurance Committee: Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. Genet Med. 2015 May;17(5):405-424
2. Tangye SG, Al-Herz W, Bousfiha A, et al: Human inborn errors of immunity: 2022 update on the classification from the international union of immunological societies expert committee. J Clin Immunol. 2022 Jun 24:1–35. doi: 10.1007/s10875-022-01289-3
3. Casanova JL, Abel L: Mechanisms of viral inflammation and disease in humans. Science. 2021 Nov 26;374(6571):1080-1086. doi: 10.1126/science.abj7965
4. Zhang Q, Bastard P; COVID Human Genetic Effort, Cobat A, Casanova JL: Human genetic and immunological determinants of critical COVID-19 pneumonia. Nature. 2022 Mar;603(7902):587-598. doi: 10.1038/s41586-022-04447-0
5. Fournier B, Latour S: Immunity to EBV as revealed by immunodeficiencies. Curr Opin Immunol. 2021 Oct;72:107-115. doi: 10.1016/j.coi.2021.04.003
6. Beziat V, Casanova JL, Jouanguy E: Human genetic and immunological dissection of papillomavirus-driven diseases: new insights into their pathogenesis. Curr Opin Virol. 2021 Dec;51:9-15. doi: 10.1016/j.coviro.2021.09.002

Performance

Method Description

Next-generation sequencing (NGS) and/or Sanger sequencing are performed to test for the presence of variants in coding regions and intron/exon boundaries of the genes analyzed, as well as some other regions that have known disease-causing variants. The human genome reference GRCh37/hg19 build was used for sequence read alignment. At least 99% of the bases are covered at a read depth over 30X. Sensitivity is estimated at above 99% for single nucleotide variants, above 94% for deletions/insertions (delins) less than 40 base pairs (bp), and above 95% for deletions up to 75 bp and insertions up to 47 bp. NGS and/or a polymerase chain reaction-based quantitative method is performed to test for the presence of deletions and duplications in the genes analyzed.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. See [Targeted Genes and Methodology Details for Viral Susceptibility, Defects in Intrinsic and Innate Immunity, Gene Panel](#) for details regarding the targeted genes analyzed for each test and specific gene regions not routinely covered.(Unpublished Mayo method)

Confirmation of select reportable variants may be performed by alternate methodologies based on internal laboratory criteria.

Genes analyzed: *CARMIL2, CD27, CD70, CIB1, CTPS1, CXCR4, DBR1, IFIH1, IFNAR1, IFNAR2, IRF3, IRF7, IRF9, MAGT1,*

POLR3A, POLR3C, PRKCD, RASGRP1, SH2D1A, STAT1, STAT2, TLR3, TLR7, TLR8, TMC6, TMC8, TNFRSF9, TRAF3, UNC93B1, and XIAP

PDF Report

Supplemental

Day(s) Performed

Varies

Report Available

28 to 42 days

Specimen Retention Time

Whole blood: 2 weeks (if available); Extracted DNA: 3 months; Cultured fibroblasts, skin biopsy: 1 month

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

81443
88233-Tissue culture, skin, solid tissue biopsy (if appropriate)
88240-Cryopreservation (if appropriate)

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
VIRID	Viral Susceptibility Gene Panel	103742-3

Result ID	Test Result Name	Result LOINC® Value
619901	Test Description	62364-5
619902	Specimen	31208-2
619903	Source	31208-2

Test Definition: VIRID

Viral Susceptibility, Defects in Intrinsic and
Innate Immunity, Gene Panel, Varies

619904	Result Summary	50397-9
619905	Result	82939-0
619906	Interpretation	69047-9
619907	Additional Results	82939-0
619908	Resources	99622-3
619909	Additional Information	48767-8
619910	Method	85069-3
619911	Genes Analyzed	82939-0
619912	Disclaimer	62364-5
619913	Released By	18771-6