
Overview**Useful For**

Determination of T-cell clonality in T-cell neoplasms

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
IHMTO	IHC Multiplex, Tech Only	No	No

Testing Algorithm

For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

Method Name

Immunohistochemistry (IHC)

NY State Available

Yes

Specimen**Specimen Type**

TECHONLY

Ordering Guidance

This test includes only technical performance of the stain; **no pathologist interpretation is provided.**

Technical component only stains **should not** be ordered with PATHC / Pathology Consultation. If ordered with PATHC, the technical component stains **will be canceled.** Any immunohistochemistry (IHC)/in situ hybridization (ISH) stain performed as a part of the PATHC will be performed at the reviewing pathologist's discretion at an additional charge.

Shipping Instructions

Attach the green "Attention Pathology" address label (T498) and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

Specimen Required

Specimen Type: Tissue

Test Definition: TRBC3

T-Cell Receptor Constant Beta Chain (TRBC1)
+CD3 Immunostain, Technical Component
Only

Supplies: Immunostain Technical Only Envelope (T693)

Container/Tube: Immunostain Technical Only Envelope

Submit:

-Formalin-fixed, paraffin-embedded tissue block

OR

-2 Unstained, positively charged glass slides (25- x 75- x 1-mm) per test ordered; sections 4-microns thick

Digital Image Access

1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: <https://news.mayocliniclabs.com/pathology/digital-imaging/>
2. Clients ordering stains using a manual requisition form will not have access to digital images.
3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: <https://news.mayocliniclabs.com/pathology/digital-imaging/#section3>

Forms

If not ordering electronically, complete, print, and send a [Immunohistochemical \(IHC\)/In Situ Hybridization \(ISH\) Stains Request](#) (T763) with the specimen.

Reject Due To

Wet/frozen tissue Cytology smears Nonformalin fixed tissue Nonparaffin embedded tissue Noncharged slides ProbeOn slides Snowcoat slides	Reject
---	--------

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive

Clinical Information

Each alpha beta T cell expresses a T-cell receptor (TCR) containing one of the two TCR beta chain constant regions (TRBC), TRBC1 or TRBC2. Reactive T-cell populations show a polytypic mixture of TRBC1 and TRBC2 expression. T-cell neoplasms show restricted expression with T-cells expressing either TRBC1 or TRBC2 in a mutually exclusive pattern, analogous to light chain expression in B-cells.

Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

The charge of glass slides can be affected by environmental factors and subsequently may alter slide staining. Sending unsuitable glass slides can result in inconsistent staining due to poor slide surface chemistry.

Best practices for storage of positively charged slides:

- Minimize time slides are stored after being unpackaged
- Limit exposure to high humidity and heat
- Minimize exposure to plastics

Avoid prolonged exposure of slides to light as this can cause the chromogens to fade over time. This fading can be mitigated by storing your slides in the dark when not being observed under the microscope.

Clinical Reference

1. Zhou T, Sardana R, Eren OC, et al. The diagnostic utility of TRBC1 immunohistochemistry in mature T-cell lymphomas. *Mod Pathol*. Published online January 28, 2025.. doi:10.1016/j.modpat.2025.100725
2. Horna P, Weybright MJ, Ferrari M, et al. Dual T-cell constant beta chain (TRBC)1 and TRBC2 staining for the identification of T-cell neoplasms by flow cytometry. *Blood Cancer J*. 2024;14(1):34
3. Soilleux EJ, Rodgers DT, Situ JJ, et al. Demonstration of T-cell monotypia using anti-TCRbeta1/2 (*TRBC1/2*) immunostaining as a rapid and cost-effective alternative to PCR-based clonality studies for the diagnosis of T-cell lymphoma. *Diagnostics (Basel)*. 2024;14(22):2479
4. Nocco SE, Ewalt MD, Moy AP, et al. TRBC1 immunohistochemistry distinguishes cutaneous T-cell lymphoma from inflammatory dermatitis: A retrospective analysis of 39 cases. *J Am Acad Dermatol*. 2024;90(4):839-841

Performance**Method Description**

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 3 days

Specimen Retention Time

Until staining is completed

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

88344-TC

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
TRBC3	TRBC1+CD3 IHC, Tech Only	No LOINC Needed

Result ID	Test Result Name	Result LOINC® Value
-----------	------------------	---------------------

Test Definition: TRBC3

T-Cell Receptor Constant Beta Chain (TRBC1)
+CD3 Immunostain, Technical Component
Only

622846	TRBC1+CD3 IHC, Tech Only	No LOINC Needed
--------	--------------------------	-----------------