

Chikungunya IgM, Antibody, Serum

## Overview

### **Useful For**

Aiding in the diagnosis of recent infection with Chikungunya virus detecting IgM antibodies in patients with recent travel to endemic areas and a compatible clinical syndrome

#### **Method Name**

Only orderable as part of a profile. For more information see CHIKV / Chikungunya IgM and IgG, Antibody, Serum.

Enzyme-Linked Immunosorbent Assay (ELISA)

### **NY State Available**

Yes

## **Specimen**

## **Specimen Type**

Serum

## **Specimen Required**

Only orderable as part of a profile. For more information see CHIKV / Chikungunya IgM and IgG, Antibody, Serum.

**Supplies:** Sarstedt Aliquot Tube, 5 mL (T914)

Collection Container/Tube:
Preferred: Serum gel
Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 0.5 mL

**Collection Instructions**: Centrifuge and aliquot serum into a plastic vial.

## **Specimen Minimum Volume**

0.4 mL

## **Reject Due To**

Gross hemolysis	Reject
Gross lipemia	Reject
Gross icterus	Reject

## **Specimen Stability Information**



Chikungunya IgM, Antibody, Serum

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	30 days	
	Frozen	30 days	

# **Clinical & Interpretive**

#### **Clinical Information**

Chikungunya virus (ChikV) is a single-stranded RNA alphavirus and a member of the Togaviridae family of viruses. The name Chikungunya is derived from the language of the Makonde ethnic groups in southeast Africa and means "that which bends" or "stooped walk." This is in reference to the hunched-over appearance of infected individuals due to the characteristically painful and incapacitating arthralgia caused by the virus. ChikV is endemic throughout Africa, India, and, more recently, the Caribbean islands. In 2014, the first case of autochthonous, or local transmission, in the United States occurred in Florida.

Humans are the primary reservoir for ChikV and *Aedes* species mosquitos are the primary vectors for transmission. Unlike other mosquito-borne viruses such as West Nile virus and Dengue, the majority of individuals who are exposed to ChikV become symptomatic, with the most severe manifestations observed at the extremes of age and in those with suppressed immunity. Once exposed to ChikV, individuals develop lasting immunity and protection from reinfection.

Prior to development of symptoms, the incubation period ranges, on average, from 3 to 7 days. Infected patients typically present with sudden-onset high fever, incapacitating joint pain, and often a maculopapular rash lasting anywhere from 3 to 10 days. Notably, symptom relapse can occur in some individuals 2 to 3 months following resolution of initial symptoms. Currently, there are no licensed vaccines and treatment is strictly supportive care.

## **Reference Values**

Only orderable as part of a profile. For more information see CHIKV / Chikungunya IgM and IgG, Antibody, Serum.

### Negative

Reference values apply to all ages.

### Interpretation

IgM and IgG Negative:

-No serologic evidence of exposure to Chikungunya virus. Repeat testing on a new specimen collected in 5 to 10 days is recommended if clinical suspicion persists.

### IgM and IgG Positive:

-IgM and IgG antibodies to Chikungunya virus detected, suggesting recent or past infection. Correlate with exposure history and clinical presentation. Confirmatory testing at a public health lab is suggested. IgM antibodies to Chikungunya virus may remain detectable for 3 to 4 months post-infection.

### IgM Positive, IgG Negative:

-IgM antibodies to Chikungunya virus detected, suggesting recent infection. Correlate with exposure history and clinical presentation. Confirmatory testing at a public health lab is suggested. Repeat testing in 5 to 10 days is recommended to



Chikungunya IgM, Antibody, Serum

demonstrate anti-Chikungunya virus IgG seroconversion to confirm current infection.

IgM Negative, IgG Positive:

-IgG antibodies to Chikungunya virus detected, suggesting past infection.

IgM and/or IgG Borderline:

-Repeat testing in 10 to 14 days is recommended.

#### **Cautions**

Specimens collected too early following infection may be negative for antibodies to Chikungunya virus. Testing of convalescent serum is recommended.

Chikungunya and Dengue viruses currently co-circulate in endemic areas and infections can present similarly in symptomatic patients. It is therefore recommended to evaluate at-risk patients for infection with both viruses.

#### **Clinical Reference**

Lwande OW, Obanda V, Bucht G, et al. Global emergence of Alphaviruses that cause arthritis in humans. Infect Ecol Epidemiol. 2015;5:29853. doi:10.3402/iee.v5.29853

### **Performance**

### **Method Description**

For the Chikungunya virus IgM assay, polystyrene microwells are coated with recombinant Chikungunya antigen. Diluted serum samples and controls are incubated in the wells to allow anti-Chikungunya antibodies (if present in the sample) to react with the antigen. Nonspecific reactants are removed by washing. Next, peroxidase-conjugated antihuman antibody is added to the wells and will react with human antibodies bound to the antigen. Excess conjugate is removed by washing. Enzyme substrate and chromogen are added, and the color is allowed to develop. After adding the Stop Reagent, the resultant color change is quantified by a spectrophotometric reading of optical density (OD). Sample OD readings are compared with reference cut-off OD readings to determine the qualitative results. (Package inserts: Anti-Chikungunya virus ELISA. IgM Euroimmun Ag;12/20/2018)

### PDF Report

No

## Day(s) Performed

Bimonthly on the second and forth Wednesday

### Report Available

Same day/1 to 14 days

### **Specimen Retention Time**

14 days

# **Performing Laboratory Location**



Chikungunya IgM, Antibody, Serum

Mayo Clinic Laboratories - Rochester Superior Drive

### **Fees & Codes**

### **Fees**

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

### **Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

### **CPT Code Information**

86790

## **LOINC®** Information

Test ID	Test Order Name	Order LOINC® Value
CHIKM	Chikungunya IgM, Ab, S	88629-1

Result ID	Test Result Name	Result LOINC® Value
CHIKM	Chikungunya IgM, Ab, S	88629-1