

# **Test Definition: FOPIA**

Opiates, Serum or Plasma, Quantitative

### **Overview**

#### **Method Name**

Quantitative Liquid Chromatography-Tandem Mass Spectrometry

### **NY State Available**

Yes

# Specimen

# **Specimen Type**

Varies

## **Specimen Required**

Submit only one of the following specimens:

Plasma

Specimen Type: Plasma

Container/Tube: Gray top (potassium oxalate/sodium fluoride), Green top (sodium heparin), Lavender top (EDTA), or

pink top (K2EDTA)

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Collection Instructions: Draw blood in a gray top (potassium oxalate/sodium fluoride), green (sodium heparin), lavender

(EDTA) or pink (K2EDTA) tube(s). Spin down and send 1 mL of plasma refrigerated in a plastic vial.

Note: Label specimen appropriately (plasma).

Serum

Specimen Type: Serum

**Collection Container/Tube:** Red top **Submission Container/Tube:** Plastic vial

Specimen Volume: 1 mL

Collection Instructions: Draw blood in a plain, red-top tube(s). Spin down and send 1 mL of serum refrigerated in a

plastic vial.

**Note:** Label specimen appropriately (serum).

## Specimen Minimum Volume

0.5 mL

### **Reject Due To**



# **Test Definition: FOPIA**

Opiates, Serum or Plasma, Quantitative

Hemolysis	Mild Reject; Gross Reject	
Lipemia	Mild OK; Gross OK	
Other	Separator tubes, Plasma or Whole blood collected in light blue (sodium citrate), specimens exposed	
	to repeat freeze/thaw cycles.	

### **Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Varies	Refrigerated (preferred)	14 days	
	Ambient	7 days	
	Frozen		

# **Clinical & Interpretive**

#### **Reference Values**

Drugs covered: codeine, morphine, 6-acetylmorphine, hydrocodone, hydromorphone, oxycodone, and oxymorphone. All drugs covered and the non-glucuronidated (free) form.

Positive cutoff: 2 ng/mL

For medical purposes only; not valid for forensic use.

### Interpretation

Identification of specific drug(s) taken by specimen donor is problematic due to common metabolites, some of which are prescription drugs themselves. The absence of expected drug(s) and/or drug metabolite(s) may indicate non-compliance, inappropriate timing of specimen collection relative to drug administration, poor drug absorption, or limitations of testing. All drugs covered are the nonglucuronidated (free) form. The concentration value must be greater than or equal to the cutoff to be reported as positive. A very small amount of an unexpected drug analyte in the presence of a large amount of an expected drug analyte may reflect pharmaceutical impurity. Interpretive questions should be directed to the laboratory.

## **Performance**

## **PDF Report**

No

#### Day(s) Performed

Monday, Wednesday and Friday

## **Report Available**



# **Test Definition: FOPIA**

Opiates, Serum or Plasma, Quantitative

3-11 days

## **Performing Laboratory Location**

**ARUP Laboratories** 

## **Fees & Codes**

#### **Fees**

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

#### **Test Classification**

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

### **CPT Code Information**

80361, 80365

#### **LOINC®** Information

Test ID	Test Order Name	Order LOINC® Value
FOPIA	Opiates, Serum or Plasma, Quant.	8217-2

Result ID	Test Result Name	Result LOINC® Value
Z4427	6-acetylmorphine, S/P, Quant	12788-6
Z4428	Codeine, S/P, Quant	3506-3
Z4429	Morphine, S/P, Quant	3827-3
Z4430	Hydrocodone, S/P, Quant	3680-6
Z4431	Hydromorphone, S/P, Quant	3683-0
Z4432	Oxycodone, S/P, Quant	3893-5
Z4433	Oxymorphone, S/P, Quant	60467-8