

Overview

Method Name

Double Immunodiffusion (DID)

NY State Available

Yes

Specimen

Specimen Type

Serum

Specimen Required

Patient Preparation: Patient should be fasting for eight hours to avoid lipemic sample interference.

Specimen Type: Serum

Container/Tube: Red top or SST

Specimen Volume: 1 mL

Collection Instructions: Draw blood in a plain red-top tube, serum gel tube(s) is also acceptable. Separate serum immediately after coagulation (30 minutes) to prevent hemolysis. Send 1 mL of serum frozen in a plastic vial.

Specimen Minimum Volume

0.5 mL

Reject Due To

Hemolysis	Reject
Lipemia	Reject
Other	Gross bacterial contamination

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated	14 days	
	Frozen (preferred)	14 days	

Clinical & Interpretive

Reference Values

Aspergillus fumigatus Neg: <1:1
Aspergillus flavus Neg: <1:1
Aspergillus niger Neg: <1:1

Performance

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

4 to 9 days

Performing Laboratory Location

LabCorp Burlington

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

CPT Code Information

86606 x 3

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
FASAB	Aspergillus Ab, Qn, DID	Not Provided

Result ID	Test Result Name	Result LOINC® Value
Z5651	Aspergillus fumigatus	23821-2
Z5652	Aspergillus flavus	23820-4
Z5653	Aspergillus niger	10894-4