

Hexosaminidase A, Serum

## **Overview**

### **Useful For**

Second-order test for diagnosing the B1 variant of Tay-Sachs disease

This test is **not useful for** testing for Sandhoff disease.

## **Testing Algorithm**

For more information see <u>Tay-Sachs and Related Disorders Diagnostic Testing Algorithm</u>

## **Special Instructions**

- Informed Consent for Genetic Testing
- Biochemical Genetics Patient Information
- Tay-Sachs and Related Disorders Diagnostic Testing Algorithm
- Informed Consent for Genetic Testing (Spanish)

#### **Method Name**

Fluorometric

#### **NY State Available**

Yes

## **Specimen**

# Specimen Type

Serum

# **Ordering Guidance**

Testing for Tay-Sachs Disease and Sandhoff Disease

The following tests are available for diagnostic and carrier testing for Tay-Sachs and Sandhoff diseases.

NAGR / Hexosaminidase A and Total, Leukocytes/Molecular Reflex, Whole Blood:

- -This is the recommended test for carrier testing for Tay-Sachs disease and Sandhoff disease.
- -Testing begins with hexosaminidase A and total enzyme analysis. If the results are consistent with an affected or carrier for Tay-Sachs disease or Sandhoff disease, next-generation sequencing to detect single nucleotide and copy number variants for *HEXA* or *HEXB*, respectively, will automatically be performed on the original specimen.
- -This test is appropriate for males and pregnant or nonpregnant females.

NAGW / Hexosaminidase A and Total Hexosaminidase, Leukocytes:

- -This test can be used for diagnosis and carrier testing for Tay-Sachs disease or Sandhoff disease.
- -Results for hexosaminidase A and total enzyme analysis are reported with recommendations for additional testing



Hexosaminidase A, Serum

when appropriate. All follow-up testing must be ordered separately on new specimens.

-This test is appropriate for males and pregnant or nonpregnant females.

## NAGS / Hexosaminidase A and Total Hexosaminidase, Serum:

- -This test can be used for diagnosis and carrier testing for Tay-Sachs disease or Sandhoff disease. Results for hexosaminidase A and total enzyme analysis are reported with recommendations for additional testing when appropriate.
- -If results indicate normal, indeterminate, or carrier status and the suspicion of Tay-Sachs disease remains high, MUGS / Hexosaminidase A, Serum for Tay-Sachs disease (B1 variant) can typically be added and performed on the same specimen.
- -With the exception of MUGS, all follow-up testing must be ordered separately on new specimens.
- -This test is **not** appropriate for pregnant females or women receiving hormonal contraception. This test is appropriate for males and nonpregnant females.
- -This test is particularly useful when it is difficult to obtain enough blood to perform leukocyte testing (NAGR or NAGW), as may be the case with infants.

### MUGS / Hexosaminidase A, Serum:

- -This is the recommended test for diagnosis and carrier testing for the B1 variant of Tay-Sachs disease. This test will not detect Sandhoff disease.
- -This test should **not** be ordered as a first-line test. Rather, this test should be ordered when the NAGR, NAGW, or NAGS indicate normal, indeterminate, or carrier results and the suspicion of Tay-Sachs disease remains high. In most cases, this test can be performed on the original specimen collected for NAGS.

#### **Necessary Information**

Healthcare professional name and phone number are required.

## Specimen Required

**Patient Preparation:** 

Fasting: 4 hours, required; Infants and small children should have specimen collected before next feeding/meal

**Supplies:** Sarstedt Aliquot Tube, 5 mL (T914)

**Collection Container/Tube:** 

Preferred: Serum gel
Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Collection Instructions: Centrifuge and aliquot serum into plastic vial.

#### **Forms**

- 1. **New York Clients-Informed consent is required.** Document on the request form or electronic order that a copy is on file. The following documents are available:
- -Informed Consent for Genetic Testing (T576)
- -Informed Consent for Genetic Testing-Spanish (T826)
- 2. Biochemical Genetics Patient Information (T602)
- 3. If not ordering electronically, complete, print, and send a <u>Biochemical Genetics Test Request</u> (T798) with the specimen.



Hexosaminidase A, Serum

# Specimen Minimum Volume

0.15 mL

## **Reject Due To**

| Gross         | OK |
|---------------|----|
| hemolysis     |    |
| Gross lipemia | OK |

## **Specimen Stability Information**

| Specimen Type | Temperature        | Time     | Special Container |
|---------------|--------------------|----------|-------------------|
| Serum         | Frozen (preferred) | 365 days |                   |
|               | Refrigerated       | 5 days   |                   |

## Clinical & Interpretive

#### **Clinical Information**

Tay-Sachs and Sandhoff diseases, also referred to as GM2 gangliosidoses, are lysosomal storage disorders caused by deficiencies of the enzymes hexosaminidase A and hexosaminidase B, respectively. These isoenzymes are dimers that differ in their subunit composition. Hexosaminidase A is a heterodimer composed of 1 alpha and 1 beta subunit (alpha-beta), while hexosaminidase B is a homodimer composed of 2 beta subunits (beta-beta). The defective lysosomal degradation and the excessive accumulation of GM2 ganglioside and related glycolipids result in the development of the clinical symptomology observed in Tay-Sachs and Sandhoff diseases.

Tay-Sachs and Sandhoff diseases are autosomal recessive conditions. Tay-Sachs disease results from 2 variants in *HEXA*, which encodes for the alpha subunit of hexosaminidase and causes a deficiency of hexosaminidase A enzyme. An increased carrier frequency for Tay-Sachs disease is observed in individuals of Ashkenazi Jewish, Celtic, and French-Canadian ancestry. Patients with Sandhoff disease have 2 variants in *HEXB*, which encodes for the beta subunit of hexosaminidase and results in deficiencies in both hexosaminidase A and hexosaminidase B enzymes. Sandhoff disease does not exhibit an increased carrier frequency in any specific population.

### Clinical Phenotypes:

Phenotypically, patients with Tay-Sachs and Sandhoff diseases are virtually indistinguishable. Variability is observed with respect to age of onset and clinical symptoms. Enzyme analysis is generally required to distinguish between the 2 disorders.

The acute infantile forms of Tay-Sachs and Sandhoff diseases typically present with progressive motor deterioration beginning at 3 to 6 months of age. Patients exhibit weakness, hypotonia, and decreasing attentiveness. Motor skills learned previously, such as crawling or sitting alone, are nearly always lost by 1 year of age. Other symptoms include rapid diminishing of vision, seizures, macrocephaly due to cerebral gliosis, and the characteristic cherry-red spot in the retina. Affected individuals typically do not survive past 5 years of age.

The juvenile or subacute forms often present between 2 and 10 years of age with ataxia and clumsiness. Patients



Hexosaminidase A, Serum

develop difficulties with speech and cognition. Neurologic features progressively get worse, and death typically occurs 2 to 4 years later.

Disease progression is slower in patients with chronic or adult-onset Tay-Sachs and Sandhoff diseases. Early signs and symptoms may be subtle and nonspecific, involving muscle and/or neurologic findings, often resulting in initial misdiagnoses. Affected individuals may exhibit abnormalities of gait and posture, spasticity, dysarthria, and progressive muscle wasting and weakness. Cognitive impairment, dementia, or psychiatric findings are observed in some patients. Significant clinical variability exists both between and within families.

#### **Testing Options:**

Several tests are available for the detection of carriers of, and individuals affected with, Tay-Sachs and Sandhoff diseases (see table below, Ordering Guidance, and Testing Algorithm). The recommended test for both diagnostic and carrier testing is NAGR / Hexosaminidase A and Total, Leukocytes/Molecular Reflex, Whole Blood. Testing begins with enzyme analysis and, when indicated, reflexes to the appropriate molecular analysis (either *HEXA* or *HEXB* gene), which includes sequencing and deletion/duplication analysis.

Follow-up molecular testing is recommended for all individuals with enzyme results in the carrier, possible carrier, or affected ranges. This differentiates between nondisease-causing pseudodeficiency alleles and disease-causing variants. In addition, molecular analysis allows for the facilitation of carrier testing and prenatal diagnosis for at-risk individuals.

| Test ID | Test Name   | Tay-Sachs<br>disease |              | Sandhoff<br>disease |              | Reflexes<br>to<br>molecular | Use during pregnancy or hormonal | Preferred<br>use                    |
|---------|---|----------------------|--------------|---------------------|--------------|-----------------------------|----------------------------------|-------------------------------------|
|         |   | Carrie<br>r          | Affecte<br>d | Carrie<br>r         | Affecte<br>d | genetic<br>testing          | contraceptio<br>n                |                                     |
| NAGR    | Hexosaminidase A<br>and Total,<br>Leukocytes/Molecul<br>ar Reflex, Whole<br>Blood | Yes                  | Yes          | Yes                 | Yes          | Yes                         | Yes                              | Diagnostic<br>or carrier<br>testing |
| NAGW    | Hexosaminidase A<br>and Total<br>Hexosaminidase,<br>Leukocytes                    | Yes                  | Yes          | Yes                 | Yes          | No                          | Yes                              | Diagnostic<br>or carrier<br>testing |
| NAGS    | Hexosaminidase A<br>and Total<br>Hexosaminidase,<br>Serum                         | Yes                  | Yes          | Yes                 | Yes          | No                          | No                               | Diagnostic                          |
| MUGS*   | Hexosaminidase A,<br>Serum  | Yes                  | Yes          | No                  | No           | No                          | No                               | Diagnostic,<br>secondary<br>only    |

<sup>\*</sup>MUGS testing should be utilized only when one of the other assays indicates normal, indeterminate, or carrier results and the clinical suspicion of Tay-Sachs disease remains high.



Hexosaminidase A, Serum

## **Reference Values**

1.23-2.59 U/L (Normal) 1.16-1.22 U/L (Indeterminate) 0.58-1.15 U/L (Carrier)

## Interpretation

Interpretation is provided with report.

### **Cautions**

GM2 activator deficiency (AB variant, *GM2A*) is a rare disorder with clinical features similar to Tay-Sachs and Sandhoff diseases; however, levels of both hexosaminidase A and B are normal. GM2 activator deficiency is not detected with this assay. Molecular genetic analysis of *GM2A* is available; see CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies.

#### Supportive Data

Regression analysis shows a linear relationship between Mayo Clinic Laboratories' MUG and 4-MUGS methods (R=0.964). A normal range of 1.23 to 2.59 U/L hexosaminidase A was determined using the 4-methylumbelliferyl-beta-D-*N*-acetyl-glucosamine-6-sulfate (4-MUGS) procedure on 50 normal subjects. A Tay-Sachs carrier range, based on results from 12 obligate carriers for Tay-Sachs disease, has been established tentatively as 0.58 to 1.15 U/L. This range will continue to be evaluated as more obligate carriers for Tay-Sachs disease are studied.

#### Clinical Reference

- 1. Tutor JC. Biochemical characterization of the GM2 gangliosidosis B1 variant. Braz J Med Biol Res. 2004;37(6):777-783
- 2. Gravel RA, Kaback MM, Proia RL, Sandhoff K, Suzuki K, Suzuki K. The GM2 gangliosidoses. In: Valle DL, Antonarakis S, Ballabio A, Beaudet AL, Mitchell GA. eds. The Online Metabolic and Molecular Bases of Inherited Disease. McGraw-Hill; 2019. Accessed September 10, 2024. Available at:

http://ommbid.mhmedical.com/content.aspx?bookid=2709&sectionid=225547784

- 3. Toro C, Shirvan L, Tifft C: *HEXA* disorders. In: Adam MP, Everman DB, Mirzaa GM, et al, eds. GeneReviews [Internet]. University of Washington, Seattle; 1999. Updated October 1, 2020. Accessed September 10, 2024. Available at: www.ncbi.nlm.nih.gov/books/NBK1218/
- 4. Leal AF, Benincore-Florez E, Solano-Galarza D, et al. GM2 Gangliosidoses: Clinical Features, Pathophysiological Aspects, and Current Therapies. Int J Mol Sci. 2020;21(17):6213. Published 2020 Aug 27. doi:10.3390/ijms21176213

#### **Performance**

#### **Method Description**

The substrate, 4-methylumbelliferyl-beta-D-*N*-acetyl-glucosamine-6-sulfate (4-MUGS), is highly specific for hexosaminidase A with minimal hydrolysis by placental hexosaminidase and none by hexosaminidase B. Hexosaminidase A hydrolyzes the 4-MUGS substrate, producing free 4-methylumbelliferyl, which fluoresces at an alkaline pH. Hexosaminidase A activity is determined by the amount of fluorescence produced.(Ben-Yoseph Y, Reid JE, Shapiro B, Nadler HL. Diagnosis and carrier detection of Tay-Sachs disease: direct determination of hexosaminidase A using 4-methylumbelliferyl derivatives of beta-N-acetylglucosamine-6-sulfate and beta-N-acetylgalactosamine-6-sulfate. Am J Hum Genet. 1985;37(4):733-740; Cowan T, Pasquali M: Laboratory Investigations of Inborn Errors of Metabolism. In: Sarafoglou K, Hoffman GF, Roth KS eds. Pediatric Endocrinology and Inborn Errors of Metabolism. 2nd ed. McGraw-Hill;



Hexosaminidase A, Serum

2017:1139-1158)

## **PDF Report**

No

## Day(s) Performed

Once per month

## **Report Available**

2 to 8 days

# **Specimen Retention Time**

30 days

# **Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

## **Fees & Codes**

#### **Fees**

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

#### **Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

## **CPT Code Information**

83080

#### **LOINC®** Information

| Test ID | Test Order Name            | Order LOINC® Value |
|---------|----------------------------|--------------------|
| MUGS    | Hexosaminidase A (MUGS), S | 2643-5             |
|         |                            |                    |

| Result ID | Test Result Name           | Result LOINC® Value |
|-----------|----------------------------|---------------------|
| 80350     | Hexosaminidase A (MUGS), S | 2643-5              |