

Overview

Useful For

Monitoring patients receiving 5-alpha reductase inhibitor therapy or chemotherapy

Evaluating patients with possible 5-alpha reductase deficiency

Testing Algorithm

For more information see [Steroid Pathways](#).

Special Instructions

- [Steroid Pathways](#)

Method Name

Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)
Portions of this test are covered by patents held by Quest Diagnostics

NY State Available

Yes

Specimen

Specimen Type

Serum

Specimen Required

Supplies: Sarstedt Aliquot Tube, 5 mL (T914)
Collection Container/Tube:
Preferred: Red top
Acceptable: Serum gel
Submission Container/Tube: Plastic vial
Specimen Volume: 1 mL
Collection Instructions: Centrifuge and aliquot serum into a plastic vial.

Specimen Minimum Volume

0.6 mL

Reject Due To

Gross hemolysis	OK
-----------------	----

Gross lipemia	OK
Gross icterus	OK

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	28 days	
	Ambient	28 days	
	Frozen	90 days	

Clinical & Interpretive

Clinical Information

The principal prostatic androgen is dihydrotestosterone (DHT). Levels of DHT remain normal with aging, despite a decrease in the plasma testosterone, and are not elevated in benign prostatic hyperplasia.(1)

Dihydrotestosterone is generated by reduction of testosterone by 5-alpha reductase. Two isoenzymes of 5-alpha reductase have been discovered. Type 1 is present in most tissues in the body where 5-alpha reductase is expressed and is the dominant form in sebaceous glands. Type 2 is the dominant isoenzyme in genital tissues, including the prostate.

Androgenetic alopecia (AGA; male-pattern baldness) is a hereditary and androgen-dependent progressive thinning of the scalp hair that follows a defined pattern.(2) While the genetic involvement is pronounced, but poorly understood, major advances have been achieved in understanding the principal elements of androgen metabolism that are involved. DHT may be related to baldness. High concentrations of 5-alpha reductase have been found in frontal scalp and genital skin and androgen-dependent processes are predominantly due to the binding of DHT to the androgen receptor. Since the clinical success of treatment of AGA with modulators of androgen metabolism or hair growth promoters is limited, sustained microscopic follicular inflammation with connective tissue remodeling, eventually resulting in permanent hair loss, is considered a possible cofactor in the complex etiology of AGA.

Available AGA treatment modalities with proven efficacy are oral finasteride, a competitive inhibitor of 5-alpha reductase type 2, and topical minoxidil, an adenosine triphosphate-sensitive potassium channel opener that has been reported to stimulate the production of vascular endothelial growth factor in cultured dermal papilla cells.

Currently, many patients with prostate disease receive treatment with a 5-alpha reductase inhibitor such as finasteride (Proscar) to diminish conversion of DHT from testosterone.

For more information see [Steroid Pathways](#).

Reference Values

Males

Cord blood: < or =100 pg/mL

< or =6 months: < or =1,200 pg/mL

Tanner Stages

Mean	Age	Reference range (pg/mL)
------	-----	-------------------------

Stage I (>6 months and prepubertal)	7.1 years	< or =50
Stage II	12.1 years	< or =200
Stage III	13.6 years	80-330
Stage IV	15.1 years	220-520
Stage V	18 years	240-650

>19 years: 112-955 pg/mL

Females

Cord blood: < or =50 pg/mL

< or =6 months: < or =1,200 pg/mL

Tanner Stages

Mean	Age	Reference range (pg/mL)
Stage I (>6 months and prepubertal)	7.1 years	< or =50
Stage II	10.5 years	< or =300
Stage III	11.6 years	< or =300
Stage IV	12.3 years	< or =300
Stage V	14.5 years	< or =300

20-55 years: < or =300 pg/mL

>55 years: < or =128 pg/mL

1. Pang S, Levine LS, Chow D, Sagiani F, Saenger P, New MI. Dihydrotestosterone and its relationship to testosterone in infancy and childhood. J Clin Endocrinol Metab. 1979;48(5):821-826

2. Stanczyk FZ. Diagnosis of hyperandrogenism: biochemical criteria. Best Pract Res Clin Endocrinol Metab. 2006;20(2):177-191

Interpretation

Patients taking 5-alpha reductase inhibitor have decreased dihydrotestosterone (DHT) serum levels.

Patients with genetic 5-alpha reductase deficiency (a rare disease) also have reduced DHT serum levels.

Dihydrotestosterone should serve as the primary marker of peripheral androgen production. However, because it is metabolized rapidly and has a very high affinity for sex hormone-binding globulin, DHT does not reflect peripheral androgen action. Instead, its distal metabolite, 3-alpha, 17-beta-androstenediol glucuronide, serves as a better marker of peripheral androgen action.

For more information see [Steroid Pathways](#).

Cautions

Patients with benign prostatic hyperplasia or prostatic cancer may not have elevated dihydrotestosterone (DHT) levels even though growth of the prostate gland may be stimulated by DHT.

Clinical Reference

1. Bartsch G, Rittmaster RS, Klocker H. Dihydrotestosterone and the concept of 5 alpha-reductase inhibition in human

benign prostatic hyperplasia. World J Urol 2002;19(6):413-425

2. Trueb RM. Molecular mechanisms of androgenetic alopecia. Exp Gerontol. 2002;37(8-9):981-990

3. Singh SM, Gauthier S, Labrie F. Androgen receptor antagonists (antiandrogens): structure-activity relationships. Curr Med Chem 2000;7(2):211-247

4. Rhodes L, Harper J, Uno H, et al. The effects of finasteride (Proscar) on hair growth, hair cycle stage, and serum testosterone and dihydrotestosterone in adult male and female stump-tail macaques (*Macaca arctoides*). J Clin Endocrinol Metab. 1994;79:991-996

5. Gustafsson O, Norming U, Gustafsson S, et al. Dihydrotestosterone and testosterone levels in men screened for prostate cancer: a study of a randomized population. Br J Urol. 1996;77:433-440

6. van der Veen A, van Faassen M, de Jong WHA, van Beek AP, Dijck-Brouwer DAJ, Kema IP. Development and validation of a LC-MS/MS method for the establishment of reference intervals and biological variation for five plasma steroid hormones. Clin Biochem. 2019;68:15-23. doi:10.1016/j.clinbiochem.2019.03.013

7. Kinter KJ, Amraei R, Anekar AA. Biochemistry, Dihydrotestosterone. In: StatPearls [Internet]. StatPearls Publishing. Last updated July 30, 2023. Accessed April 14, 2025. Available at www.ncbi.nlm.nih.gov/books/NBK557634

Performance

Method Description

The assay uses high performance liquid chromatography with triple quadrupole tandem mass spectrometry. Deuterated stable isotope of dihydrotestosterone (13C3-DHT) is added to a 0.5 mL serum sample as internal standard. The DHT and internal standard are extracted from the sample by liquid/liquid extraction. This is followed by high performance liquid chromatography on a Cohesive LX4 System and analysis on a tandem mass spectrometer equipped with an electrospray ion source.(Unpublished Mayo Method)

PDF Report

No

Day(s) Performed

Monday, Wednesday, Friday

Report Available

2 to 8 days

Specimen Retention Time

2 weeks

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Superior Drive

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

82642

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
DHTS	Dihydrotestosterone, S	1848-1

Result ID	Test Result Name	Result LOINC® Value
81479	Dihydrotestosterone, S	1848-1