

## Overview

### Useful For

Diagnosing and monitoring liver disease, particularly diseases resulting in a destruction of hepatocytes

### Method Name

Photometric Rate, L-Aspartate with Pyridoxyl-5-Phosphate

### NY State Available

Yes

## Specimen

### Specimen Type

Serum

### Necessary Information

Patient's age and sex are required.

### Specimen Required

#### Collection Container/Tube:

**Preferred:** Serum gel

**Acceptable:** Red top

**Specimen Volume:** 0.5 mL

**Submission Container/Tube:** Plastic vial

#### Collection Instructions:

1. Serum gel tubes should be centrifuged within 2 hours of collection.
2. Red-top tubes should be centrifuged, and the serum aliquoted into a plastic vial within 2 hours of collection.

### Forms

If not ordering electronically, complete, print, and send a [Kidney Transplant Test Request](#) with the specimen.

### Specimen Minimum Volume

0.25 mL

### Reject Due To

Gross hemolysis	Reject
-----------------	--------

**Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	7 days	
	Ambient	7 days	
	Frozen	30 days	

**Clinical & Interpretive****Clinical Information**

Aspartate aminotransferase (AST) is found in high concentrations in liver, heart, skeletal muscle, and kidney. AST is present in both cytoplasm and mitochondria of cells. In cases involving mild tissue injury, the predominant form of AST is that from the cytoplasm. Severe tissue damage results in more of the mitochondrial enzyme being released. High levels of AST can be found in cases such as myocardial infarction, acute liver cell damage, viral hepatitis, and carbon tetrachloride poisoning. Slight to moderate elevation of AST is seen in muscular dystrophy, dermatomyositis, acute pancreatitis, and crushed muscle injuries.

**Reference Values**

Males

0-11 months: not established

1-13 years: 8-60 U/L

&gt; or =14 years: 8-48 U/L

Females

0-11 months: not established

1-13 years: 8-50 U/L

&gt; or =14 years: 8-43 U/L

**Interpretation**

Elevated aspartate aminotransferase (AST) values are seen in parenchymal liver diseases characterized by a destruction of hepatocytes. Values are typically at least 10 times above the normal range. Levels may reach values as high as 100 times the upper reference limit, although 20- to 50-fold elevations are most frequently encountered. In infectious hepatitis and other inflammatory conditions affecting the liver, alanine aminotransferase (ALT) is characteristically as high as or higher than AST, and the ALT:AST ratio, which normally and in other condition is less than 1, becomes greater than unity. AST levels are usually elevated before clinical signs and symptoms of disease appear. Five- to 10-fold elevations of both AST and ALT occur in patients with primary or metastatic carcinoma of the liver, with AST usually being higher than ALT, but levels are often normal in the early stages of malignant infiltration of the liver. Elevations of ALT activity persist longer than do those of AST activity. Elevated AST values may also be seen in disorders affecting the heart, skeletal muscle, and kidney.

**Cautions**

Pyridoxal phosphate is a cofactor in the reaction and is necessary for enzyme activity.

**Clinical Reference**

Tietz Textbook of Clinical Chemistry. Edited by CA Burtis, ER Ashwood. Philadelphia, WB Saunders Company, 1994

## Performance

### Method Description

Aspartate aminotransferase (AST) is measured by a coupled enzyme kinetic method where the rate of decrease of NADH, determined at 340 nm, is directly proportional to the AST activity. (Package insert: Roche AST reagent, Indianapolis, IN, January 2000)

### PDF Report

No

### Day(s) Performed

Monday through Sunday

### Report Available

Same day/1 to 2 days

### Specimen Retention Time

1 week

### Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

### Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

### CPT Code Information

84450

### LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
---------	-----------------	--------------------

## Test Definition: AST

Aspartate Aminotransferase (AST) (GOT),  
Serum

AST	Aspartate Aminotransferase (AST), S	30239-8
-----	-------------------------------------	---------

Result ID	Test Result Name	Result LOINC® Value
AST	Aspartate Aminotransferase (AST), S	30239-8