

# **Test Definition: CL**

Chloride, Serum

### Overview

**Useful For** Evaluation of water, electrolyte, and acid-base status

Method Name Potentiometric/Indirect Ion-Selective Electrode

NY State Available Yes

Specimen

Specimen Type Serum

**Necessary Information** Patient's age and sex are required.

Specimen Required
Collection Container/Tube:
Preferred: Serum gel
Acceptable: Red top
Submission Container/Tube: Plastic vial
Specimen Volume: 0.5 mL
Collection Instructions:
1. Serum gel tubes should be centrifuged within 2 hours of collection.
2. Red-top tubes should be centrifuged, and the serum aliquoted into a plastic vial within 2 hours of collection.

### Forms

If not ordering electronically, complete, print, and send a <u>Renal Diagnostics Test Request</u> (T830) with the specimen.

#### Specimen Minimum Volume

0.25 mL

### **Reject Due To**

Gross	Reject
hemolysis	

### **Specimen Stability Information**

# **Test Definition: CL**



Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	7 days	
	Frozen	30 days	

## Clinical & Interpretive

## **Clinical Information**

Chloride is the major anion in the extracellular water space; its physiological significance is in maintaining proper body water distribution, osmotic pressure, and normal anion-cation balance in the extracellular fluid compartment.

Chloride is increased in dehydration, renal tubular acidosis (hyperchloremia metabolic acidosis), acute renal failure, metabolic acidosis associated with prolonged diarrhea and loss of sodium bicarbonate, diabetes insipidus, adrenocortical hyperfunction, salicylate intoxication, and with excessive infusion of isotonic saline or extremely high dietary intake of salt. Hyperchloremia acidosis may be a sign of severe renal tubular pathology.

Chloride is decreased in overhydration, chronic respiratory acidosis, salt-losing nephritis, metabolic alkalosis, congestive heart failure, Addisonian crisis, certain types of metabolic acidosis, persistent gastric secretion and prolonged vomiting, aldosteronism, bromide intoxication, syndrome of inappropriate antidiuretic hormone secretion, and conditions associated with expansion of extracellular fluid volume.

## **Reference Values**

1-17 years: 102-112 mmol/L
> or =18 years: 98-107 mmol/L
Reference values have not been established for patients who are under 12 months of age.

### Interpretation

In normal individuals, serum chloride values vary little during the day, although there is a slight decrease after meals due to the diversion of chloride to the production of gastric juice.

### Cautions

High serum values of other halide ions may lead to falsely high readings on the chloride ion-selective electrode.

### **Clinical Reference**

Tietz Textbook of Clinical Chemistry. Edited by CA Burtis, ER Ashwood. WB Saunders Company, Philadelphia, PA, 1994

## Performance

### **Method Description**

The ion-selective electrode (ISE) module of the P-Module performs indirect measurement of electromotive force (EMF). The ISE module measures the EMF difference between an ISE and a reference electrode. The EMF of the ISE is dependent on the ion concentration of the sample. The EMF of the reference electrode is constant. The P-Module uses an electronic calculation circuit to convert EMF of the sample to the ion concentration of the sample.(Package insert: Roche Diagnostics ISE reagent, Indianapolis, IN, 2006)



Chloride, Serum

## PDF Report

No

# Day(s) Performed

Monday through Sunday

# Report Available

Same day/1 to 2 days

## **Specimen Retention Time**

1 week

## **Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

## Fees & Codes

### Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

### **Test Classification**

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

## **CPT Code Information**

82435

## LOINC<sup>®</sup> Information

Test ID	Test Order Name	Order LOINC <sup>®</sup> Value
CL	Chloride, S	2075-0
02		20/0 0

Result ID	Test Result Name	Result LOINC <sup>®</sup> Value
CL	Chloride, S	2075-0