

Overview

Useful For

Diagnosing deficiency of coagulation factor XI

Investigating prolonged activated partial thromboplastin time

Special Instructions

- [Coagulation Guidelines for Specimen Handling and Processing](#)

Method Name

Optical Clot-Based

NY State Available

Yes

Specimen

Specimen Type

Plasma Na Cit

Ordering Guidance

Coagulation testing is highly complex, often requiring the performance of multiple assays and correlation with clinical information. For that reason, consider ordering a Coagulation Consultation.

Necessary Information

If priority specimen, mark request form, give reason, and request a call-back.

Specimen Required

Specimen Type: Platelet-poor plasma

Patient Preparation: Patient must not be receiving Coumadin (warfarin) or heparin therapy.

Collection Container/Tube: Light-blue top (3.2% sodium citrate)

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Collection Instructions:

1. Specimen must be collected prior to factor replacement therapy.
2. For complete instructions, see [Coagulation Guidelines for Specimen Handling and Processing](#).
3. Centrifuge, transfer all plasma into a plastic vial, and centrifuge plasma again.
4. Aliquot plasma into a plastic vial, leaving 0.25 mL in the bottom of centrifuged vial.
5. Freeze plasma immediately (no longer than 4 hours after collection) at -20 degrees C or ideally, at or below -40 degrees C.

Additional Information:

1. Double-centrifuged specimen is critical for accurate results as platelet contamination may cause spurious results.
2. Each coagulation assay requested should have its own vial.

Forms

[If not ordering electronically, complete, print, and send a Coagulation Test Request](#) (T753) with the specimen.

Specimen Minimum Volume

0.5 mL

Reject Due To

| | |
|-----------------|--------|
| Gross hemolysis | Reject |
| Gross lipemia | Reject |
| Gross icterus | Reject |

Specimen Stability Information

| Specimen Type | Temperature | Time | Special Container |
|---------------|-------------|---------|-------------------|
| Plasma Na Cit | Frozen | 14 days | |

Clinical & Interpretive

Clinical Information

Factor XI is synthesized in the liver. Its biological half-life is 60 to 80 hours. Factor XI is a component of intrinsic coagulation pathway and, when activated, activates factor IX to IXa.

Factor XI deficiency may cause prolonged partial thromboplastin time. Deficiency is associated with mild bleeding diathesis, but there is poor correlation between activity level and clinical bleeding. A relatively high incidence of congenital deficiency occurs among individuals of Ashkenazi Jewish descent (hemophilia C).

Reference Values

Adults: 55-150%

Normal, full-term newborn infants or healthy premature infants may have decreased levels (> or =10%), which may not reach adult levels [for 180 or more days postnatal.*](#)

*See Pediatric Hemostasis References section in [Coagulation Guidelines for Specimen Handling and Processing](#).

Interpretation

Acquired deficiency is associated with liver disease and rarely inhibitors.

Patients who are homozygous: <20% activity

Patients who are heterozygous: 20% to 60% activity

Cautions

Decreased plasma levels of factor XI do not correlate well with bleeding risk.

Clinical Reference

1. He R, Chen D, He S. Factor XI: hemostasis, thrombosis, and antithrombosis. *Thromb Res.* 2012;129(5):541-550
2. Martin-Salces M, Jimenez-Yuste V, Alvarez MT, Quintana M, Hernandez-Navarro F. Review: Factor XI deficiency: review and management in pregnant women. *Clin Appl Thromb Hemost.* 2010;16(2):209-213
3. Seligsohn U. Factor XI in haemostasis and thrombosis: past, present and future. *Thromb Haemost.* 2007;98(1):84-89
4. Santoro R, Prejano S, Iannaccaro P. Factor XI deficiency: a description of 34 cases and literature review. *Blood Coagul Fibrinolysis.* 2011;22(5):431-435
5. Favaloro EJ, Lippi G, eds. *Hemostasis and Thrombosis: Methods and Protocols.* Humana Press; 2017

Performance

Method Description

The factor XI assay is performed on the Instrumentation Laboratory ACL TOP using the activated partial thromboplastin time (aPTT) method and a factor-deficient substrate. Patient plasma is combined and incubated with a factor XI-deficient substrate (normal plasma depleted of factor XI by immunoadsorption) and an aPTT reagent. After a specified incubation time, calcium is added to trigger the coagulation process in the mixture. Then the time to clot formation is measured optically at a wavelength of 671 nm.(Owen CA Jr, Bowie EJW, Thompson JH Jr. *Diagnosis of Bleeding Disorders.* 2nd ed. Little, Brown and Company; 1975; Cielsa B. Defects of plasma clotting factors. In: *Hematology in Practice.* 3rd ed. FA Davis; 2019:chap 17)

PDF Report

No

Day(s) Performed

Monday through Saturday

Report Available

1 to 3 days

Specimen Retention Time

7 days

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.

- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been modified from the manufacturer's instructions. Its performance characteristics were determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

85270

LOINC® Information

| Test ID | Test Order Name | Order LOINC® Value |
|---------|-------------------------|--------------------|
| F_11 | Coag Factor XI Assay, P | 3226-8 |

| Result ID | Test Result Name | Result LOINC® Value |
|-----------|-------------------------|---------------------|
| F_11 | Coag Factor XI Assay, P | 3226-8 |