

**Overview****Method Name**

Enzyme-Linked Immunosorbent Assay (ELISA)

**NY State Available**

No

**Specimen****Specimen Type**

Serum

**Specimen Required**

**Specimen Type:** Serum

**Container/Tube:** Red Top or SST

**Specimen Volume:** 3 mL

**Collection Instructions:** Draw blood in a plain red-top tube(s), serum gel tube is acceptable. Spin down and send 3 mL of serum frozen in a plastic vial.

**Specimen Minimum Volume**

0.5 mL

**Reject Due To**

Gross hemolysis	Reject
Gross lipemia	Reject
List other reasons for rejection	Microbial contaminants

**Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Serum	Frozen (preferred)	30 days	
	Refrigerated	14 days	

**Clinical & Interpretive**

**Clinical Information**

The anti-phospholipid syndrome (APS) is a disorder of recurrent vascular thrombosis associated with persistently positive anticardiolipin (aCL) or lupus anticoagulant tests. In patients with APS, anticardiolipin antibodies bind a variety of charged phospholipids, including phosphatidylethanolamine, as well as they do cardiolipin. Lupus patients also have high titers of autoantibodies to various phospholipids, including phosphatidylethanolamine.

Presentations of the syndrome include thrombosis of deep veins of the legs, as well as renal, hepatic, inferior vena cava or sagittal veins. Occlusion of the arterial circulation may be manifested as stroke, ischemic retinopathy, myocardial or bowel infarction, or peripheral gangrene. Thrombosis can occur in veins or arteries of any size. Recurrent pregnancy loss also appears to be the result of thrombosis within the placental vasculature.

Anti-phospholipid antibody tests are supplemental tests and should not be used alone for diagnostic purposes. Diagnosis of anti-phospholipid syndrome must be made in conjunction with other clinical indications.

**Reference Values**

Anti-Phosphatidylcholine IgA: <12.0 U/mL

Anti-Phosphatidylcholine IgG: <12.0 U/mL

Anti-Phosphatidylcholine IgM: <12.0 U/mL

Reference Range applies to Antiphosphatidylcholine IgA, IgG & IgM:

Normal <12.0

Equivocal 12.0 – 18.0

Elevated >18.0

**Performance****PDF Report**

No

**Day(s) Performed**

Wednesday

**Report Available**

3 to 11 days

**Performing Laboratory Location**

BioAgilytix Diagnostics

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

### Test Classification

The performance characteristics of the listed assay was validated by BioAgilytix Diagnostics. The US FDA has not approved or cleared this test. The results of this assay can be used for clinical diagnosis without FDA approval.

BioAgilytix Diagnostics is a CLIA certified, CAP accredited laboratory for performing high complexity assays such as this one.

### CPT Code Information

83520/x3

### LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
FCLNE	Anti-Phosphatidylcholine Panel	Not Provided

Result ID	Test Result Name	Result LOINC® Value
Z0149	Anti-phosphatidylcholine IgG	13073-2
Z0140	Anti-Phosphatidylcholine IgM	13074-0
Z0141	Anti-Phosphatidylcholine IgA	13075-7