
Overview

Useful For

Detection of acute hepatitis C virus (HCV) infection before the appearance of HCV antibodies in serum (ie, <2 months from exposure)

Detection and confirmation of chronic HCV infection

Quantification of HCV RNA in serum of patients with chronic HCV infection (HCV antibody-positive)

Monitoring disease progression in chronic HCV infection and response to antiviral therapy

Determining cure and detection of relapse after completion of antiviral therapy

Testing Algorithm

For information see:

[-Chronic Hepatitis C Treatment and Monitoring Algorithm: Direct Antiviral Antigen \(DAA\) Combination](#)

[-Hepatitis C: Testing Algorithm for Screening and Diagnosis](#)

Special Instructions

- [Hepatitis C: Testing Algorithm for Screening and Diagnosis](#)
- [Chronic Hepatitis C Treatment and Monitoring Algorithm: Direct Antiviral Agent \(DAA\) Combination](#)

Highlights

A reflex test for serum specimens that are hepatitis C virus (HCV) antibody screen-reactive for diagnosis of chronic hepatitis C.

This test is appropriate for diagnosis of acute hepatitis C in high-risk or immunosuppressed individuals who may be negative for HCV antibodies.

This test can be used to establish a baseline HCV viral load before initiating antiviral therapy for chronic hepatitis C.

This test can be used to monitor response and progress during antiviral therapy for chronic hepatitis C.

This test is appropriate for confirming a sustained virologic response and detecting a relapse of hepatitis C after completion of antiviral therapy.

Method Name

Real-Time Reverse Transcription-Polymerase Chain Reaction (RT-PCR)

NY State Available

Yes

Specimen**Specimen Type**

Serum SST

Ordering Guidance

For detection and quantification of hepatitis C viral RNA in serum before, during, and after antiviral therapy for chronic hepatitis C.

Shipping Instructions

1. Freeze serum immediately, and ship specimen frozen on dry ice only.
2. If shipment will be delayed for more than 24 hours, freeze serum at -20 to -80 degrees C (up to 84 days) until shipment on dry ice.

Specimen Required**Supplies:** Sarstedt 5 mL Aliquot Tube (T914)**Collection Container/Tube:** Serum gel**Submission Container/Tube:** Plastic vial**Specimen Volume:** 1.5 mL Serum**Collection Instructions:** Within 2 hours of collection, centrifuge and aliquot serum into a plastic vial.**Forms**

If not ordering electronically, complete, print, and send 1 of the following forms with the specimen:

[-Gastroenterology and Hepatology Test Request \(T728\)](#)

[-Microbiology Test Request \(T244\)](#)

[-General Test Request \(T239\)](#)

[-Kidney Transplant Test Request](#)

Specimen Minimum Volume

Serum: 0.8 mL

Reject Due To

Gross hemolysis	Reject
Gross lipemia	OK
Gross icterus	OK

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum SST	Frozen (preferred)	84 days	
	Refrigerated	6 days	

Clinical & Interpretive**Clinical Information**

About 75% of all individuals infected with hepatitis C virus (HCV) will develop chronic hepatitis C with ongoing viral replication in the liver and detectable HCV RNA in serum or plasma, eventually resulting in cirrhosis. The remaining 25% of the infected individuals recover from the infection without evidence of viral replication or presence of detectable HCV RNA in serum or plasma. Chronic HCV infection can be cured at variable success rates with either combined interferon-alpha and ribavirin therapy or interferon-free combination of direct-acting antiviral (DAA) agents.

The antiviral response rates correlate with pretreatment serum or plasma HCV RNA levels (viral load) and the HCV genotype found in the infected individuals. The optimal duration of combined interferon and ribavirin therapy can be determined from the patient's pretreatment viral load and HCV genotype. Clinical trial studies indicated that a decrease in HCV RNA levels of more than 2 log IU/mL at 4 weeks or 12 weeks of therapy is predictive of an increased chance of achieving a sustained virologic response (defined as undetectable HCV RNA levels in serum 6 months after completing antiviral therapy). Despite receiving longer duration of antiviral therapy (48 weeks versus 24 weeks), patients with chronic infection due to HCV genotypes 1 and 4 generally have less favorable sustained virologic response rates (40%-50%) than those infected with genotypes 2 and 3 (>80%). Due to the necessary prolonged duration (typically 24- to 48-week duration) and low cure rates of such antiviral therapy, interferon-based therapy has been supplanted with potent interferon-free DAA combination therapy.

Cure rates, as defined by sustained virologic response, of over 90% are observed among patients infected with HCV and treated with interferon-free DAA combinations that are of shorter treatment duration (eg, 8 or 12 weeks) than those of interferon-based therapy. Current guidelines for antiviral therapy of chronic hepatitis C recommend quantitative testing for HCV RNA in serum or plasma before initiating antiviral therapy, at 4 weeks of therapy, and 12 weeks after completion of therapy. A HCV RNA level below 25 IU/mL in serum or plasma 12 weeks after ending therapy is the therapeutic goal and indicates an SVR is achieved. Quantitative HCV RNA testing can be considered at the end of therapy and at 24 weeks or later after completion of antiviral therapy.

For more information, see the following:

[-Chronic Hepatitis C Treatment and Monitoring Algorithm: Direct Antiviral Antigen \(DAA\) Combination](#)

[-Hepatitis C: Testing Algorithm for Screening and Diagnosis](#)

Reference Values

Undetected

Interpretation

This assay has a result range of 15 to 100,000,000 IU/mL (1.18 log to 8.00 log IU/mL) for quantification of hepatitis C virus (HCV) RNA in serum.

An "Undetected" result indicates that the HCV is absent in the patient's serum specimen.

A result of "<15 IU/mL (<1.18 log IU/mL)" indicates that HCV RNA is detected, but the HCV RNA level present cannot be quantified accurately below this lower limit of quantification of this assay. When clinically indicated, follow-up testing with this assay is recommended in 1 to 2 months. To assess response-guided therapy eligibility, an "Undetected" result is required, and a result of "<15 IU/mL (<1.18 log IU/mL)" should not be considered equivalent to an "Undetected" result.

A quantitative result expressed in IU/mL and log IU/mL indicates the degree of active HCV viral replication in the patient. Monitoring HCV RNA levels over time is important to assess disease progression and/or monitoring a patient's response to anti-HCV therapy.

A result of ">100,000,000 IU/mL (>8.00 log IU/mL)" indicates the presence of active HCV viral replication, and the HCV RNA level present cannot be quantified accurately above this upper limit of quantification of this assay.

An "Inconclusive" result reported with a comment indicates that testing failed, likely due to presence of inhibitory substances in the submitted serum specimen. A new specimen should be collected for retesting.

Cautions

Except for immunocompromised patients or patients with suspected acute hepatitis, laboratory evaluation of hepatitis C virus (HCV) infection status should begin with HCV serologic testing, including testing for the presence of HCV antibodies (see [Hepatitis C: Testing Algorithm for Screening and Diagnosis](#)). A diagnosis of chronic HCV infection should not be based solely on the presence of detectable or quantifiable HCV RNA in a single serum specimen.

An "Undetected" HCV RNA test result with a "Reactive" HCV antibody screen result may be due to:

1. A false-reactive HCV antibody screen result
2. Resolved or past HCV infection
3. Transient low viremia (ie, episodic viral replication) of active HCV infection

To distinguish between the first 2 conditions, another HCV antibody test (eg, HCCAD / Hepatitis C Virus Antibody Screen for Cadaveric or Hemolyzed Specimens, Serum) can be requested. To distinguish between the latter 2 conditions, patients should be retested for HCV RNA in 1 to 2 months, as clinically indicated.

Clinical Reference

1. de Leuw P, Sarrazin C, Zeuzem S. How to use virological tools for the optimal management of chronic hepatitis C. *Liver Int.* 2011;31 Suppl 1:3-12
2. Centers for Disease Control and Prevention (CDC): Testing for HCV infection: an update of guidance for clinicians and laboratorians. *MMWR Morb Mortal Wkly Rep.* 2013;62(18):362-365
3. American Association for the Study of Liver Diseases, Infectious Diseases Society of America: HCV guidance: Recommendations for testing, managing, and treating hepatitis C. AASLD, IDSA; Accessed April 29, 2026. Available at www.hcvguidelines.org/contents

Performance**Method Description**

The cobas hepatitis C virus (HCV) assay is a US Food and Drug Administration-approved, in vitro nucleic acid amplification test for the quantification of HCV RNA in human serum using the cobas x800 Systems for fully automated viral nucleic acid extraction (generic silica-based capture technique) and automated amplification and detection of the viral nucleic acid sequence. This polymerase chain reaction (PCR) assay amplifies sequences within the highly conserved 5' noncoding region of the HCV genome and generates amplification products that are detected and quantified in real-time with 2 sequence-specific TaqMan probes. A non-HCV armored RNA quantitation standard (RNA QS) is introduced into each specimen during sample preparation to serve as internal control for nucleic acid extraction and PCR amplification/detection processes. Fluorescent reporter dye-labeled TaqMan probes hybridized to the complementary HCV target sequences and RNA QS sequence undergo hydrolysis during PCR amplification step to generate fluorescent signal detected in 2 different dye channels. Concentration of the HCV RNA in a patient's serum sample is determined by a ratio of the intensity of the fluorescent dye from the cleaved HCV target sequence probes and that from the RNA QS target probe detected throughout the PCR process. (Package insert: cobas HCV-Quantitative nucleic acid test for use on the cobas 5800/6800/8800 Systems. Roche Molecular Systems, Inc; Rev. 6.0, 11/2025)

PDF Report

No

Day(s) Performed

Monday through Saturday

Report Available

1 to 3 days

Specimen Retention Time

2 months

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Superior Drive

Fees & Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Definition: HCVQN

Hepatitis C Virus (HCV) RNA Detection and
Quantification by Real-Time Reverse
Transcription-PCR, Serum

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

87522

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
HCVQN	HCV RNA Detect/Quant, S	11011-4

Result ID	Test Result Name	Result LOINC® Value
97291	HCV RNA Detect/Quant, S	11011-4